

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

P

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07C 251/60, A01N 37/00</b>		<b>A1</b>	(11) International Publication Number: <b>WO 95/18789</b>
			(43) International Publication Date: <b>13 July 1995 (13.07.95)</b>
(21) International Application Number: <b>PCT/EP94/04318</b>		(81) Designated States: AM, AU, BB, BG, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, NO, NZ, PL, RO, RU, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).	
(22) International Filing Date: <b>27 December 1994 (27.12.94)</b>			
(30) Priority Data: 12/94-9                      5 January 1994 (05.01.94)                      CH 2117/94-0                      1 July 1994 (01.07.94)                      CH			
(71) Applicant (for all designated States except US): <b>CIBA-GEIGY AG [CH/CH]; Klybeckstrasse 141, CH-4002 Basle (CH).</b>		<b>Published</b> <i>With international search report.</i>	
(72) Inventors; and			
(75) Inventors/Applicants (for US only): <b>ZIEGLER, Hugo [CH/CH]; Im Bohnacker 15, CH-4108 Witterswil (CH). TRAH, Stephan [DE/DE]; Am Hertweg 14 A, D-79110 Freiburg (DE). FAROOQ, Saleem [CH/CH]; Kirchackerstrasse 27, CH-4422 Arisdorf (CH). ZURFLÜH, René [CH/CH]; St. Alban Rheinweg 180, CH-4052 Basle (CH).</b>			
(74) Common Representative: <b>CIBA-GEIGY AG; Patentabteilung, Klybeckstrasse 141, CH-4002 Basle (CH).</b>			

(54) Title: PESTICIDES



(57) Abstract

Compounds of formula (I) and the isomers, and isomer mixtures thereof, which are possible, in which either a) X is an N atom and Y is OR<sub>11</sub> or N(R<sub>12</sub>)R<sub>13</sub>, or b) X is CH and Y is OR<sub>11</sub>, and in which furthermore: R<sub>11</sub> is C<sub>1</sub>-C<sub>4</sub>alkyl; R<sub>12</sub> and R<sub>13</sub>, independently, are hydrogen or C<sub>1</sub>-C<sub>4</sub>alkyl; A is an O atom or the group NR<sub>4</sub>; R<sub>1</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, halo-C<sub>1</sub>-C<sub>4</sub>alkyl, cyclopropyl, cyano or methylthio; R<sub>2</sub> is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, a group (i), a group (ii), or thienyl; R<sub>3</sub> is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>haloalkyl having 1 to 5 halogen atoms, C<sub>1</sub>-C<sub>4</sub>alkoxy-C<sub>1</sub>-C<sub>2</sub>alkyl, C<sub>2</sub>-C<sub>4</sub>alkenyl-C<sub>1</sub>-C<sub>2</sub>alkyl, which is unsubstituted or substituted by 1 to 3 halogen atoms, C<sub>2</sub>-C<sub>4</sub>alkynyl-C<sub>1</sub>-C<sub>2</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl which is unsubstituted or substituted by 1 to 4 halogen atoms, C<sub>3</sub>-C<sub>6</sub>cycloalkyl-C<sub>1</sub>-C<sub>4</sub>alkyl which is unsubstituted or substituted by 1 to 4 halogen atoms, cyano-C<sub>1</sub>-C<sub>4</sub>alkyl; C<sub>1</sub>-C<sub>4</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>2</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxycarbamoyl-C<sub>1</sub>-C<sub>2</sub>alkyl, phenyl-C<sub>1</sub>-C<sub>3</sub>alkyl which is unsubstituted or substituted by halogen, C<sub>1</sub>-C<sub>3</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>haloalkyl, cyano, nitro or C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, it being possible for the phenyl group to be monosubstituted to trisubstituted by identical or different substituents; phenyl which is unsubstituted or mono- to disubstituted, independently, by C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, halogen, C<sub>1</sub>-C<sub>2</sub>haloalkyl having 1 to 3 halogen atoms, nitro or cyano, or pyridyl which is unsubstituted or mono- to disubstituted, independently, by C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, halogen, C<sub>1</sub>-C<sub>2</sub>haloalkyl having 1 to 3 halogen atoms, nitro or cyano; R<sub>4</sub> is C<sub>1</sub>-C<sub>4</sub>alkyl, phenyl, or R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 7-membered ring which is unsubstituted or substituted by C<sub>1</sub>-C<sub>4</sub>alkyl and which can have 1 to 3 additional hetero atoms selected from amongst N, O and S, and in which the substituents Z, B, D and n are as defined in the present publication, are pesticidal active ingredients. They can be used for pest control, in particular as microbicides, insecticides and acaricides in agriculture, horticulture and in the hygiene sector.

BEST AVAILABLE COPY


**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Pesticides

The invention relates to novel pesticidally active compounds of the formula I



and to the isomers, and isomer mixtures, thereof which are possible, in which either

- a) X is an N atom and  
Y is OR<sub>11</sub> or N(R<sub>12</sub>)R<sub>13</sub>, or
- b) X is CH and  
Y is OR<sub>11</sub>,


and in which furthermore:


R<sub>11</sub> is C<sub>1</sub>-C<sub>4</sub>alkyl;

R<sub>12</sub> and R<sub>13</sub>, independently, are hydrogen or C<sub>1</sub>-C<sub>4</sub>alkyl;

A is an O atom or the group NR<sub>4</sub>;

R<sub>1</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, halo-C<sub>1</sub>-C<sub>4</sub>alkyl, cyclopropyl, cyano or methylthio;

R<sub>2</sub> is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, a group ,

a group  Z-B, or thienyl;

D radicals are identical or different and are halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy,

C<sub>1</sub>-C<sub>2</sub>haloalkyl, C<sub>1</sub>-C<sub>2</sub>haloalkoxy, C<sub>3</sub>-C<sub>6</sub>alkenyloxy, C<sub>3</sub>-C<sub>6</sub>alkynyloxy,

C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, cyano or nitro;

n is 0, 1, 2, 3 or 4;

Z is -O-, -O-(C<sub>1</sub>-C<sub>4</sub>alkyl)-, -(C<sub>1</sub>-C<sub>4</sub>alkyl)-O-, -S(O)<sub>m</sub>-, -(C<sub>1</sub>-C<sub>4</sub>alkyl)-S(O)<sub>m</sub>-,

-S(O)<sub>m</sub>-(C<sub>1</sub>-C<sub>4</sub>alkyl)-,

m is 0, 1 or 2,

B is C<sub>1</sub>-C<sub>6</sub>alkyl, halo-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, or is C<sub>2</sub>-C<sub>6</sub>alkenyl or

C<sub>2</sub>-C<sub>4</sub>alkynyl-C<sub>1</sub>-C<sub>2</sub>alkyl, each of which is unsubstituted or substituted by 1 to 3 halogen

- 2 -

atoms, or is aryl or heterocyclyl, each of these two, independently, being unsubstituted or mono- to pentasubstituted by C<sub>1</sub>-C<sub>6</sub>alkyl, halo-C<sub>1</sub>-C<sub>6</sub>alkyl, halogen, C<sub>1</sub>-C<sub>6</sub>alkoxy or

halo-C<sub>1</sub>-C<sub>6</sub>alkoxy, or a group  $\text{---}(\text{CH})_p \begin{array}{c} \text{R}_5 \quad \text{R}_6 \quad \text{R}_7 \\ \diagdown \quad \diagup \quad \diagup \\ \text{C} \\ \diagup \quad \diagdown \quad \diagdown \\ \text{R}_9 \quad \text{R}_8 \end{array}$ , or trimethylsilyl;

R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub> and R<sub>9</sub>, independently, are hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl or halogen and p is 0, 1, 2 or 3;

R<sub>3</sub> is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>haloalkyl having 1 to 5 halogen atoms, C<sub>1</sub>-C<sub>4</sub>alkoxy-C<sub>1</sub>-C<sub>2</sub>alkyl, C<sub>2</sub>-C<sub>4</sub>alkenyl-C<sub>1</sub>-C<sub>2</sub>alkyl, which is unsubstituted or substituted by 1 to 3 halogen atoms, C<sub>2</sub>-C<sub>4</sub>alkynyl-C<sub>1</sub>-C<sub>2</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl which is unsubstituted or substituted by 1 to 4 halogen atoms, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl-C<sub>1</sub>-C<sub>4</sub>alkyl which is unsubstituted or substituted by 1 to 4 halogen atoms, cyano-C<sub>1</sub>-C<sub>4</sub>alkyl; C<sub>1</sub>-C<sub>4</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>2</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>2</sub>alkyl, phenyl-C<sub>1</sub>-C<sub>3</sub>alkyl which is unsubstituted or substituted by halogen, C<sub>1</sub>-C<sub>3</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>haloalkyl, cyano, nitro or C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, it being possible for the phenyl group to be monosubstituted to trisubstituted by identical or different substituents; phenyl which is unsubstituted or mono- to disubstituted, independently, by C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, halogen, C<sub>1</sub>-C<sub>2</sub>haloalkyl having 1 to 3 halogen atoms, nitro or cyano, or pyridyl which is unsubstituted or mono- to disubstituted, independently, by C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, halogen, C<sub>1</sub>-C<sub>2</sub>haloalkyl having 1 to 3 halogen atoms, nitro or cyano;

R<sub>4</sub> is C<sub>1</sub>-C<sub>4</sub>alkyl, phenyl, or

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 7-membered ring which is unsubstituted or substituted by C<sub>1</sub>-C<sub>4</sub>alkyl and which can have 1 to 3 additional hetero atoms selected from amongst N, O and S.

The compounds according to the invention have fungicidal, acaricidal and insecticidal properties and are suitable as active ingredients for use in agriculture, horticulture and the hygiene field.

The invention furthermore also relates to a process for the preparation of the compounds according to the invention, and to fungicidal, acaricidal and insecticidal compositions which comprise such compounds as active ingredients, and to the use of such compounds and compositions for controlling phytopathogenic fungi, Acarina and insects and for preventing such an attack.

If asymmetric carbon atoms exist in the compounds of the formula I, the compounds exist

in optically active form. The compounds exist in any case in [E] and/or [Z] forms, merely because of the presence of the aliphatic, the oximino and the hydrazono double bonds. Furthermore, atropisomerism may exist. The formula I is intended to embrace all these isomeric forms which are possible and their mixtures, for example racemic mixtures, and any [E/Z] mixtures.

Unless otherwise defined, the general terms used hereinabove and hereinbelow are intended to mean the following:

Halogen is fluorine, chlorine, bromine or iodine, in particular fluorine, chlorine or bromine, especially fluorine or chlorine.

Alkyl is either straight-chain, for example methyl, ethyl, n-propyl, n-butyl, n-hexyl, n-octyl, n-decyl, n-dodecyl, n-hexadecyl or n-octadecyl, or branched, for example isopropyl, isobutyl, sec-butyl, tert-butyl, isopentyl, neopentyl or isohexyl.

Alkenyl is straight-chain or branched alkenyl, for example vinyl, 1-methylvinyl, allyl, 1-butenyl, isopropenyl, in particular allyl.

Alkynyl is, for example, ethynyl, 1-propynyl or 1-butylnyl, in particular propargyl.

Cycloalkyl is to be understood as meaning cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.

Halogen-substituted groups such as haloalkyl and haloalkoxy can be partially or fully halogenated by identical or different substituents. Examples of haloalkyl are methyl which is mono- to trisubstituted by fluorine, chlorine and/or bromine, such as  $\text{CHF}_2$ ,  $\text{CF}_3$  or  $\text{CH}_2\text{Cl}$ ; ethyl which is mono- to pentasubstituted by fluorine, chlorine and/or bromine, such as  $\text{CH}_2\text{CF}_3$ ,  $\text{CF}_2\text{CF}_3$ ,  $\text{CF}_2\text{CCl}_3$ ,  $\text{CF}_2\text{CHCl}_2$ ,  $\text{CF}_2\text{CHF}_2$ ,  $\text{CF}_2\text{CFCl}_2$ ,  $\text{CH}_2\text{CH}_2\text{Cl}$ ,  $\text{CF}_2\text{CHBr}_2$ ,  $\text{CF}_2\text{CHClF}$ ,  $\text{CF}_2\text{CHBrF}$  or  $\text{CClFCHClF}$ ; trifluoromethyl is very particularly preferred.

Straight-chain  $\text{C}_1$ - $\text{C}_4$ alkylenedioxy is  $-\text{O}-\text{CH}_2-\text{O}-$ ,  $-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-$ ,  $-\text{O}-\text{CH}_2\text{CH}_2\text{CH}_2-\text{O}-$  or  $-\text{O}-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-\text{O}-$ .

Aryl is, for example, phenyl or naphthyl, in particular phenyl.

Heterocyclyl is a 5- to 7-membered aromatic or non-aromatic ring having one to three hetero atoms selected from the group consisting of N, O and S. Preferred are aromatic 5- and 6-membered rings which have a nitrogen atom as hetero atom and, if appropriate, a further hetero atom, preferably nitrogen or sulfur, in particular nitrogen.

The term of the 5- to 7-membered ring which is formed by R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded encompasses, in particular, pyrrolidine, piperidine, morpholine, thiomorpholine, hexamethyleneimine, imidazole, pyrazole, pyrrole, 1,2,4-triazole, 1,2,3-triazole, tetrazole, isoxazole, oxazole, isoxazolidine, oxazolidine, thiazole, isothiazole, thiazoline and isothiazolidine.

Preferred within the scope of the invention are


(1) Compounds of the formula I in which either


- a) X is an N atom and  
Y is OCH<sub>3</sub> or NHCH<sub>3</sub>, or
- b) X is CH and  
Y is OCH<sub>3</sub>,

and in which furthermore:

A is an O atom or the group NR<sub>4</sub>;

R<sub>1</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, halo-C<sub>1</sub>-C<sub>4</sub>alkyl, cyclopropyl, cyano or methylthio;

R<sub>2</sub> is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, a group , a group



D radicals are identical or different and are halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>2</sub>haloalkyl, C<sub>1</sub>-C<sub>2</sub>haloalkoxy, C<sub>3</sub>-C<sub>6</sub>alkenyloxy, C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, cyano or nitro;

n is 0, 1, 2, 3 or 4;

Z is -O-, -O-(C<sub>1</sub>-C<sub>4</sub>alkyl)-, -(C<sub>1</sub>-C<sub>4</sub>alkyl)-O-, -S(O)<sub>m</sub>-, -(C<sub>1</sub>-C<sub>4</sub>alkyl)-S(O)<sub>m</sub>-, -S(O)<sub>m</sub>-(C<sub>1</sub>-C<sub>4</sub>alkyl)-,

m is 0, 1 or 2,


B is C<sub>1</sub>-C<sub>6</sub>alkyl, halo-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl,

or is C<sub>2</sub>-C<sub>6</sub>alkenyl or C<sub>2</sub>-C<sub>4</sub>alkynyl-C<sub>1</sub>-C<sub>2</sub>alkyl, each of which is unsubstituted or

- 6 -


(6) compounds of the formula I in which  $R_1$  is hydrogen, methyl, cyclopropyl or cyano, in particular methyl;

(7) compounds of the formula I in which  $R_2$  is  $C_1$ - $C_4$ alkyl or cyclopropyl, in particular methyl or cyclopropyl;

(8) compounds of the formula I in which  $R_2$  is a group  and

$D$  is halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ alkoxy,  $C_1$ - $C_2$ alkyl which is substituted by 1 to 5 halogen atoms,  $C_1$ - $C_2$ haloalkoxy,  $C_3$ - $C_6$ alkenyloxy,  $C_3$ - $C_6$ alkynyloxy,  $C_1$ - $C_4$ alkylenedioxy, cyano, or nitro, or thienyl,


$D$  is, in particular, fluorine, chlorine, bromine,  $C_1$ - $C_4$ alkyl or  $-CF_3$ ;

(9) compounds of the formula I in which  $R_2$  is a group  and


$Z$  is  $-O-$ ,  $-O-(C_1-C_4alkyl)-$ ,  $-(C_1-C_4alkyl)-O-$ ,  $-S(O)_2-$ ,  $-(C_1-C_4alkyl)-S(O)_2-$ ,  $-S(O)_2-(C_1-C_4alkyl)-$ ,

in particular  $-O-$ ,  $-CH_2-O-$  or  $-O-CH_2-$ ,

very particularly  $-O-CH_2$ ;


(10) compounds of the formula I in which  $R_2$  is a group  and

$B$  is  $C_1$ - $C_4$ alkyl, halogen- $C_1$ - $C_4$ alkyl, or is  $C_2$ - $C_4$ alkenyl or  $C_2$ - $C_4$ alkynyl- $C_1$ - $C_2$ alkyl, each of which is unsubstituted or substituted by 1 to 3 halogen atoms, or is aryl or aryl which is monosubstituted or disubstituted, independently, by  $C_1$ - $C_2$ alkyl, halo- $C_1$ - $C_2$ alkyl, halogen,

$C_1$ - $C_2$ alkoxy or halo- $C_1$ - $C_2$ alkoxy, or is a group 

$R_5$ ,  $R_6$ ,  $R_7$ ,  $R_8$  and  $R_9$ , independently, are hydrogen,  $C_1$ - $C_2$ alkyl or halogen, and  $p$  is 0, 1, 2 or 3;

$B$  is, in particular,  $C_1$ - $C_2$ alkyl, halo- $C_1$ - $C_3$ alkyl, or is allyl or propargyl, each of which is unsubstituted or in each case substituted by 1 or 2 halogen atoms or 1 or two methyl

groups, or phenyl, phenyl which is substituted by a substituent selected from the group consisting of fluorine, chlorine, bromine and  $\text{CF}_3$ , or a group  $-(\text{CH})_2$   ;

(11) compounds of the formula I in which  $\text{R}_2$  is a phenyl group which is substituted in the 4-position by -Z-B;

(12) compounds of the formula I in which

$\text{R}_3$  is hydrogen,  $\text{C}_1$ - $\text{C}_6$ alkyl or  $\text{C}_1$ - $\text{C}_4$ haloalkyl having 1 to 3 halogen atoms,  $\text{C}_1$ - $\text{C}_2$ alkoxy- $\text{C}_1$ - $\text{C}_2$ alkyl,  $\text{C}_1$ - $\text{C}_4$ alkoxycarbonyl- $\text{C}_1$ - $\text{C}_2$ alkyl, propenyl which is unsubstituted or substituted by 1 to 3 halogen atoms, propargyl,  $\text{C}_3$ - $\text{C}_6$ cycloalkyl, cyclopropylmethyl which is unsubstituted or substituted by 1 to 2 halogen atoms, cyano- $\text{C}_1$ - $\text{C}_2$ alkyl, phenyl- $\text{C}_1$ - $\text{C}_2$ alkyl which is unsubstituted or substituted by halogen, methyl, methoxy or halomethyl having 1 to 3 halogen atoms, it being possible for the phenyl group to be monosubstituted to disubstituted by identical or different substituents; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or pyridyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or  $\text{R}_3$  and  $\text{R}_4$  together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 7-membered ring which is unsubstituted or substituted by  $\text{C}_1$ - $\text{C}_4$ alkyl and which can have 1 to 3 additional hetero atoms selected from amongst N, O and S;  $\text{R}_3$  is preferably hydrogen,  $\text{C}_1$ - $\text{C}_6$ alkyl,  $\text{C}_1$ - $\text{C}_4$ haloalkyl having 1 to 3 halogen atoms,  $\text{C}_1$ - $\text{C}_2$ alkoxymethyl, prop-2-en-1-yl which is unsubstituted or substituted by 1 to 3 halogen atoms, propargyl,  $\text{C}_3$ - $\text{C}_6$ cycloalkyl, cyclopropylmethyl which is unsubstituted or substituted by 1 to 2 fluorine or chlorine atoms, cyano- $\text{C}_1$ - $\text{C}_2$ alkyl, phenyl- $\text{C}_1$ - $\text{C}_2$ alkyl which is unsubstituted or substituted by halogen, methyl, methoxy or halomethyl having 1 to 3 halogen atoms, it being possible for the phenyl group to be monosubstituted to disubstituted by identical or different substituents; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or pyridyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or  $\text{R}_3$  and  $\text{R}_4$  together with the nitrogen atom to which they are bonded are 1,2,4-triazolyl, 4-morpholinyl, 1-azepinyl, 1-piperidinyl or 1-pyrrolidinyl,  $\text{R}_3$  is particularly preferably methyl;



(13) compounds of the formula I in which  $R_4$  is methyl or phenyl, particularly methyl;

(14) compounds of the formula I in which

X is CH; Y is  $OCH_3$ ;  $R_1$  is  $CH_3$ ; A is oxygen;

$R_2$  is either 4-methylphenyl or 4-allyloxyphenyl or 4-(3-trifluoromethyl-benzyloxy)phenyl or 4-(2,2-dichlorocyclopropylmethoxy)phenyl and

$R_3$  is  $CH_3$ ;

(15) compounds of the formula I in which either

a) X is an N atom and  
Y is  $OCH_3$  or  $NHCH_3$ , or

b) X is CH and  
Y is  $OCH_3$ ,

and in which furthermore

A is an O atom or the group  $NR_4$ ;

$R_1$  is hydrogen;  $C_1$ - $C_4$ alkyl; cyclopropyl; cyano or methylthio;

$R_2$  is hydrogen;  $C_1$ - $C_6$ alkyl;  $C_3$ - $C_6$ cycloalkyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ alkoxy,  $C_1$ - $C_2$ haloalkyl,  $C_1$ - $C_2$ haloalkoxy,  $C_3$ - $C_6$ alkenyloxy,  $C_3$ - $C_6$ alkynyloxy,  $C_1$ - $C_4$ alkylenedioxy, cyano or nitro; thienyl;

$R_3$  is hydrogen;  $C_1$ - $C_6$ alkyl;  $C_1$ - $C_6$ haloalkyl having 1 to 5 halogen atoms;

$C_1$ - $C_4$ alkoxy- $C_1$ - $C_2$ alkyl;  $C_2$ - $C_4$ alkenyl- $C_1$ - $C_2$ alkyl which is unsubstituted or substituted by 1 to 3 halogen atoms;  $C_2$ - $C_4$ alkynyl- $C_1$ - $C_2$ alkyl;  $C_3$ - $C_6$ cycloalkyl which is unsubstituted or substituted by 1 to 4 halogen atoms;  $C_3$ - $C_6$ cycloalkyl- $C_1$ - $C_4$ alkyl which is unsubstituted or substituted by 1 to 4 halogen atoms; cyano- $C_1$ - $C_4$ alkyl;

$C_1$ - $C_4$ alkoxycarbonyl- $C_1$ - $C_2$ alkyl; phenyl- $C_1$ - $C_3$ alkyl which is unsubstituted or substituted by halogen,  $C_1$ - $C_3$ alkyl,  $C_1$ - $C_4$ alkoxy,  $C_1$ - $C_4$ haloalkyl, cyano, nitro,  $C_1$ - $C_4$ alkylenedioxy, it being possible for the phenyl group to be monosubstituted to trisubstituted by identical or different substituents; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ alkoxy, halogen,  $C_1$ - $C_2$ haloalkyl having 1 to 3 halogen atoms, nitro or cyano; pyridyl which is unsubstituted or monosubstituted to disubstituted, independently, by  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ alkoxy, halogen,  $C_1$ - $C_2$ haloalkyl having 1 to 3 halogen atoms, nitro or cyano;

$R_4$  is  $C_1$ - $C_4$ alkyl; phenyl; or

$R_3$  and  $R_4$  together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 7-membered ring which is unsubstituted or substituted by  $C_1$ - $C_4$ alkyl and

which can have 1 to 3 additional hetero atoms selected from amongst N, O and S.

(16) compounds of the formula I in which

X is CH or N

Y is OCH<sub>3</sub>

A is O or N-R<sub>4</sub>

R<sub>1</sub> is methyl, cyclopropyl or methylthio;

R<sub>2</sub> is methyl; cyclopropyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>2</sub>alkyl which is substituted by 1 to 5 halogen atoms, C<sub>1</sub>-C<sub>2</sub>haloalkoxy, C<sub>3</sub>-C<sub>6</sub>alkenyloxy,

C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, cyano, or nitro; or thienyl; and in which

R<sub>3</sub> is as defined for formula I and

R<sub>4</sub> is methyl or phenyl, or

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded are either pyrrolidine, piperidine, morpholine, thiomorpholine, hexamethyleneimine, imidazole, pyrazole, pyrrole, 1,2,4-triazole or 1,2,3-triazole.

(17) Compounds of the formula I in which:

X is N,

Y is NHCH<sub>3</sub>,

A is O or N-R<sub>4</sub>,

R<sub>1</sub> is methyl, cyclopropyl or methylthio;

R<sub>2</sub> is methyl; cyclopropyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>2</sub>alkyl which is substituted by 1 to 5 halogen atoms, C<sub>1</sub>-C<sub>2</sub>haloalkoxy, C<sub>3</sub>-C<sub>6</sub>alkenyloxy,

C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, cyano or nitro; or thienyl; and in which

R<sub>3</sub> is as defined for formula I and

R<sub>4</sub> is methyl or phenyl, or

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded are either pyrrolidine, piperidine, morpholine, thiomorpholine, hexamethyleneimine, imidazole, pyrazole, pyrrole, 1,2,4-triazole or 1,2,3-triazole.

(18) Compounds of the formula I in which:

A is oxygen,

R<sub>1</sub> is methyl,

R<sub>2</sub> is methyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>2</sub>alkyl which is substituted by 1

- 10 -

to 5 halogen atoms, C<sub>1</sub>-C<sub>2</sub>haloalkoxy, C<sub>3</sub>-C<sub>6</sub>alkenyloxy, C<sub>3</sub>-C<sub>6</sub>alkynyloxy,

C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, cyano or nitro; or thienyl; and

R<sub>3</sub> is C<sub>1</sub>-C<sub>6</sub>alkyl;

while X and Y are as defined for formula I.

(19) Compounds of the formula I in which

R<sub>1</sub> is methyl,

R<sub>2</sub> is methyl and

R<sub>3</sub> is as described and

R<sub>4</sub> is methyl or phenyl, or

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 7-membered ring which is unsubstituted or substituted by C<sub>1</sub>-C<sub>4</sub>alkyl and which can have 1 to 3 additional hetero atoms selected from amongst N, O and S; while A, X and Y are as defined for formula I.

(20) Compounds of the formula I in which:

R<sub>3</sub> is hydrogen; C<sub>1</sub>-C<sub>4</sub>alkyl; C<sub>1</sub>-C<sub>4</sub>haloalkyl having 1 to 3 halogen atoms;

C<sub>1</sub>-C<sub>2</sub>alkoxy-C<sub>1</sub>-C<sub>2</sub>alkyl; propenyl which is unsubstituted or substituted by 1 to 3 halogen atoms; propargyl; C<sub>3</sub>-C<sub>6</sub>cycloalkyl; C<sub>3</sub>-C<sub>6</sub>cycloalkylmethyl which is unsubstituted or substituted by 1 to 2 halogen atoms; cyano-C<sub>1</sub>-C<sub>2</sub>-alkyl;

C<sub>1</sub>-C<sub>2</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>2</sub>alkyl; phenyl-C<sub>1</sub>-C<sub>2</sub>alkyl which is unsubstituted or substituted by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano, nitro or C<sub>1</sub>-C<sub>2</sub>alkylenedioxy, it being possible for the phenyl group to be monosubstituted to disubstituted by identical or different substituents; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or pyridyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro;

R<sub>4</sub> is methyl or phenyl; or

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 7-membered ring which is unsubstituted or substituted by C<sub>1</sub>-C<sub>4</sub>alkyl and which can have 1 to 3 additional hetero atoms selected from amongst N, O and S; while A, X, Y, R<sub>1</sub> and R<sub>2</sub> are as defined for formula I.

(21) Amongst the compounds of the formula I mentioned under (20), those in which R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded are triazolyl, morpholinyl, 2,6-dimethylmorpholinyl, azepinyl, piperidyl or pyrrolidinyl.

- 11 -

(22) Compounds of the formula I in which:

A is oxygen,

R<sub>1</sub> is methyl,

R<sub>2</sub> is methyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>2</sub>alkyl which is substituted by 1 to 5 halogen atoms, C<sub>1</sub>-C<sub>2</sub>haloalkoxy, C<sub>3</sub>-C<sub>6</sub>alkenyloxy, C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, cyano or nitro; or thienyl; and

R<sub>3</sub> is hydrogen; C<sub>1</sub>-C<sub>4</sub>alkyl; C<sub>1</sub>-C<sub>4</sub>haloalkyl having 1 to 3 halogen atoms;

C<sub>1</sub>-C<sub>2</sub>alkoxy-C<sub>1</sub>-C<sub>2</sub>alkyl; propenyl which is unsubstituted or substituted by 1 to 3 halogen atoms; propargyl; C<sub>3</sub>-C<sub>6</sub>cycloalkyl; C<sub>3</sub>-C<sub>6</sub>cycloalkylmethyl which is unsubstituted or substituted by 1 to 2 halogen atoms; cyano-C<sub>1</sub>-C<sub>2</sub>-alkyl;

C<sub>1</sub>-C<sub>2</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>2</sub>alkyl; phenyl-C<sub>1</sub>-C<sub>2</sub>alkyl which is unsubstituted or substituted by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano, nitro or C<sub>1</sub>-C<sub>2</sub>alkylenedioxy, it being possible for the phenyl group to be monosubstituted to disubstituted by identical or different substituents; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or pyridyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro;

while X and Y are as defined for formula I.

(23) Amongst the compounds of the formula I mentioned under (22), those in which:

A is oxygen,

R<sub>1</sub> is methyl,

R<sub>2</sub> is methyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, trifluoromethyl or trifluoromethoxy; and

R<sub>3</sub> is methyl.

(24) Compounds of the formula I in which:

A is NCH<sub>3</sub>,

R<sub>1</sub> is methyl,

R<sub>2</sub> is methyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, trifluoromethyl or trifluoromethoxy;

R<sub>3</sub> is methyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or pyridyl which is unsubstituted or monosubstituted to disubstituted,

- 12 -

independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro;  
while X and Y are as defined for formula I.

(25) Compounds of the formula I in which:

X is an N atom;

Y is OR<sub>11</sub>;

R<sub>11</sub> is C<sub>1</sub>-C<sub>4</sub>alkyl; and in which

A, R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are as defined for formula I.

(26) Amongst the compounds mentioned under (25), those in which:

A is an O atom;

R<sub>1</sub> and R<sub>2</sub> are methyl;

R<sub>3</sub> is hydrogen; C<sub>1</sub>-C<sub>4</sub>alkyl; C<sub>1</sub>-C<sub>4</sub>haloalkyl having 1 to 3 halogen atoms;

C<sub>1</sub>-C<sub>2</sub>alkoxy-C<sub>1</sub>-C<sub>2</sub>alkyl; propenyl which is unsubstituted or substituted by 1 to 3 halogen atoms; propargyl; C<sub>3</sub>-C<sub>6</sub>cycloalkyl; C<sub>3</sub>-C<sub>6</sub>cycloalkylmethyl which is unsubstituted or substituted by 1 to 2 halogen atoms; cyano-C<sub>1</sub>-C<sub>2</sub>-alkyl;

C<sub>1</sub>-C<sub>2</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>2</sub>alkyl; phenyl-C<sub>1</sub>-C<sub>2</sub>alkyl which is unsubstituted or substituted by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano, nitro or C<sub>1</sub>-C<sub>2</sub>alkylenedioxy, it being possible for the phenyl group to be monosubstituted to disubstituted by identical or different substituents; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or pyridyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro.

(27) Amongst the compounds mentioned under (26), those in which:

R<sub>3</sub> is hydrogen; C<sub>1</sub>-C<sub>4</sub>alkyl or C<sub>1</sub>-C<sub>4</sub>haloalkyl having 1 to 3 halogen atoms.

(28) Compounds of the formula I in which:

X is an N atom;

Y is N(R<sub>12</sub>)R<sub>13</sub>;

R<sub>12</sub> and R<sub>13</sub>, independently, are hydrogen or C<sub>1</sub>-C<sub>4</sub>alkyl; and in which

A, R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are as defined for formula I.

(29) Amongst the compounds mentioned under (28), those in which:

Y is NH<sub>2</sub>, N(CH<sub>3</sub>)<sub>2</sub> or NHC<sub>2</sub>H<sub>5</sub>;

- 13 -

A is an O atom;

R<sub>1</sub> and R<sub>2</sub> are methyl;

R<sub>3</sub> is hydrogen; C<sub>1</sub>-C<sub>4</sub>alkyl; C<sub>1</sub>-C<sub>4</sub>haloalkyl having 1 to 3 halogen atoms;

C<sub>1</sub>-C<sub>2</sub>alkoxy-C<sub>1</sub>-C<sub>2</sub>alkyl; propenyl which is unsubstituted or substituted by 1 to 3 halogen atoms; propargyl; C<sub>3</sub>-C<sub>6</sub>cycloalkyl; C<sub>3</sub>-C<sub>6</sub>cycloalkylmethyl which is unsubstituted or substituted by 1 to 2 halogen atoms; cyano-C<sub>1</sub>-C<sub>2</sub>-alkyl;

C<sub>1</sub>-C<sub>2</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>2</sub>alkyl; phenyl-C<sub>1</sub>-C<sub>2</sub>alkyl which is unsubstituted or substituted by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano, nitro or C<sub>1</sub>-C<sub>2</sub>alkylenedioxy, it being possible for the phenyl group to be monosubstituted to disubstituted by identical or different substituents; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or pyridyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro.

(30) Amongst the compounds mentioned under (29), those in which:

R<sub>3</sub> is hydrogen; C<sub>1</sub>-C<sub>4</sub>alkyl or C<sub>1</sub>-C<sub>4</sub>haloalkyl having 1 to 3 halogen atoms.

(31) Compounds of the formula I in which the X=C double bond is in the E form.


The compounds of the formula I can be prepared as follows:

A) To obtain a compound of the formula I in which Y is N(R<sub>12</sub>)R<sub>13</sub>, a compound of the formula I in which Y is OR<sub>11</sub> is reacted with HN(R<sub>12</sub>)R<sub>13</sub>. The reaction is carried out advantageously in an inert organic diluent, for example in an alcohol such as ethanol, in an ether such as tetrahydrofuran or dioxane, an ester such as ethyl acetate, a sulfoxide such as dimethyl sulfoxide, an amide such as dimethylformamide or a ketone such as methyl isobutyl ketone. Methylamine can be used in gaseous form or else in dissolved form, for example in the form of a solution in ethanol. The process temperature is conventionally between 0°C and 40°C, preferably room temperature.

B) To prepare a compound of the formula I in which X, Y, A and R<sub>1</sub>-R<sub>3</sub> are as defined for formula I (and in which R<sub>3</sub> is not hydrogen):

A compound of the general formula

substituted by 1 to 3 halogen atoms, or is aryl or heterocyclyl, or aryl or heterocyclyl, each of these, independently, being monosubstituted to pentasubstituted by C<sub>1</sub>-C<sub>6</sub>alkyl, halo-C<sub>1</sub>-C<sub>6</sub>alkyl, halogen, C<sub>1</sub>-C<sub>6</sub>alkoxy or halo-C<sub>1</sub>-C<sub>6</sub>alkoxy, or a group



R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub> and R<sub>9</sub>, independently, are hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl or halogen and p is 0, 1, 2 or 3;

R<sub>3</sub> is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>haloalkyl having 1 to 5 halogen atoms, C<sub>1</sub>-C<sub>4</sub>alkoxy-C<sub>1</sub>-C<sub>2</sub>alkyl, C<sub>2</sub>-C<sub>4</sub>alkenyl-C<sub>1</sub>-C<sub>2</sub>alkyl which is unsubstituted or substituted by 1 to 3 halogen atoms, C<sub>2</sub>-C<sub>4</sub>alkynyl-C<sub>1</sub>-C<sub>2</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl which is unsubstituted or substituted by 1 to 4 halogen atoms, C<sub>3</sub>-C<sub>6</sub>cycloalkyl-C<sub>1</sub>-C<sub>4</sub>alkyl which is unsubstituted or substituted by 1 to 4 halogen atoms, cyano-C<sub>1</sub>-C<sub>4</sub>alkyl; C<sub>1</sub>-C<sub>4</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>2</sub>alkyl, phenyl-C<sub>1</sub>-C<sub>3</sub>alkyl which is unsubstituted or substituted by halogen, C<sub>1</sub>-C<sub>3</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>haloalkyl, cyano, nitro, C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, it being possible for the phenyl group to be monosubstituted to trisubstituted by identical or different substituents; phenyl which is unsubstituted or mono- to disubstituted, independently, by C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, halogen, C<sub>1</sub>-C<sub>2</sub>haloalkyl having 1 to 3 halogen atoms, nitro or cyano, or pyridyl which is unsubstituted or mono- to disubstituted, independently, by C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, halogen, C<sub>1</sub>-C<sub>2</sub>haloalkyl having 1 to 3 halogen atoms, nitro or cyano;

R<sub>4</sub> is C<sub>1</sub>-C<sub>4</sub>alkyl or phenyl, or

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 7-membered ring which is unsubstituted or substituted by C<sub>1</sub>-C<sub>4</sub>alkyl and which can have 1 to 3 additional hetero atoms selected from amongst N, O and S;


(2) compounds of the formula I in which X is N and Y is OCH<sub>3</sub>;

(3) compounds of the formula I in which X is CH;

(4) compounds of the formula I in which X is N, Y is NHCH<sub>3</sub> and R<sub>1</sub> is H, CH<sub>3</sub>, cyclopropyl or CN;


(5) compounds of the formula I in which A is oxygen, NCH<sub>3</sub> or n-C<sub>6</sub>H<sub>5</sub>, in particular oxygen or NCH<sub>3</sub>, very particularly oxygen;

- 14 -



II

in which A and R<sub>1</sub>-R<sub>3</sub> are as defined above is reacted with a compound of the general formula



III

in which X and Y are as defined above and U is a leaving group.

This reaction is a nucleophilic substitution reaction which can be carried out under the reaction conditions conventionally used in such a case. The leaving group U is preferably chlorine, bromine, iodine, mesyloxy or tosyloxy. The reaction is advantageously carried out in an inert organic diluent such as a cyclic ether, for example tetrahydrofuran or dioxane, a ketone such as acetone, an amide such as dimethylformamide, a sulfoxide such as dimethyl sulfoxide, in the presence of a base such as sodium hydride, sodium carbonate, potassium carbonate, sodium amide, a tertiary amine, for example trialkylamine, in particular diazabicyclononane or diazabicycloundecane, or silver oxide, at temperatures between -20°C and 80°C, preferably in a temperature range of 0°C to 50°C.

Alternatively, the reaction can be carried out with phase transfer catalysis in an organic solvent for example methylene chloride, in the presence of an aqueous alkaline solution, for example sodium hydroxide solution, and of a phase transfer catalyst, for example tetrabutylammonium hydrogen sulfate, at room temperature.

The resulting compounds of the formula I can be isolated and purified by methods known per se. Equally, isomer mixtures obtained, for example E/Z isomer mixtures, can be separated by methods known per se to give the pure isomers, for example by means of chromatography or fractional crystallization.




The oximes of the general formula II which are used as starting materials are either known or can be prepared by known methods (J. Chem. Soc., Perkin Trans II 537 (1990); Ber. Deutsch. Chem. Ges. 62, 866 (1929); Gazz. Chim. Ital. 37 II, 147 (1907); Liebigs Ann. Chem. 262, 305 (1891)).

Equally, the starting materials of the formula III can be prepared in a manner known per se, for example as described in the European Patent Publication EP-A-203 606 (BASF) and the literature cited therein, or as described in Angew. Chem. 71, 349-365 (1959).

C) To prepare a compound of the formula I in which A is oxygen and X, Y and R<sub>1</sub>-R<sub>3</sub> are as defined for formula I:

A compound of the general formula



IV

in which X, Y, R<sub>1</sub> and R<sub>2</sub> are as defined above is reacted with a compound of the general formula




V,

R<sub>3</sub> being as defined under formula I and U being as defined under formula III (and R<sub>3</sub> being neither hydrogen nor phenyl nor pyridyl).

This reaction is a nucleophilic substitution reaction as described under B).

D) To prepare a compound of the formula IV in which X, Y, R<sub>1</sub> and R<sub>2</sub> are as defined for formula I, a procedure is followed in which:

A compound of the general formula




VI

in which X, Y, R<sub>1</sub> and R<sub>2</sub> are as defined above is reacted with hydroxylamine or a salt thereof, for example the hydrochloride. The reaction is carried out advantageously in pyridine or methanol as the solvent at temperatures between -20°C and +80°C or the boiling point of methanol, preferably in a temperature range of 0°C to 50°C, a base being required if methanol is used, for example an alkali metal carbonate, such as potassium carbonate, a tertiary amine such as triethylamine or diazabicyclononane, pyridine or silver oxide.

The ketone of the general formula VI is prepared analogously to the method described under B). The ketones of the general formula VI and their preparation are described, for example, in EP-370 629, EP-506 149, EP-403 618, EP-414 153, EP-463 488, EP-472 300, EP-460 575, WO-92/18494 and in other publications.

E) To prepare a compound of the formula I in which A, X, Y and R<sub>1</sub>-R<sub>3</sub> are as defined for formula I but in which R<sub>3</sub> is not hydrogen, a procedure is followed in which, for example:

A compound of the general formula




VII

in which A, X, Y and R<sub>1</sub>-R<sub>3</sub> are as defined above is reacted with a methylating agent, for example methyl iodide, dimethyl sulfate or diazomethane. The reaction is advantageously carried out in the presence of a base, for example potassium carbonate or sodium hydride, in a suitable solvent and at suitable reaction temperatures (see, for example, H.S. Anker und H.T. Clarke; Organic Synthesis, Coll. Vol. 3, 172).

F) To prepare a compound of the formula VII in which A, X, Y and R<sub>1</sub>-R<sub>3</sub> are as defined for formula I, a procedure is advantageously followed in which:

A compound of the formula




VIII

in which A, Y and R<sub>1</sub>-R<sub>3</sub> are as defined above is reacted with a formate in the presence of a base analogously to the method described in EP-A-178 826, or is subjected to nitrosation with nitrous acid or a nitrite in the presence of a base analogously to the method described in EP-A-254 426.

A further possibility of synthesizing a compound of the formula VII is the following reaction:

A compound of the formula



IX

in which A, Y and R<sub>1</sub> to R<sub>3</sub> are as defined above is reacted with methoxymethylenetriphenylphosphorane analogously to the method described in EP-A-178 826 or with O-methylhydroxylamine (or a salt thereof) analogously to the method described in EP-A-254 426.

The novel compounds of the formulae VII, VIII and IX are also provided by the invention.

It has now been found that compounds of the formula I have a microbicidal spectrum for

- 18 -

controlling phytopathogenic microorganisms, in particular fungi, which is particularly favourable for practical requirements. They have very advantageous curative, preventive and, in particular, systemic properties and can be used for protecting a large number of crop plants. Using the active ingredients of the formula I, the pests which are found on plants or parts of plants (fruits, flowers, foliage, stalks, tubers, roots) of various crops of useful plants can be contained or destroyed, and even parts of plants which are formed at a later point in time remain free from phytopathogenic microorganisms.

The compounds of the formula I can furthermore be used as seed-dressing agents for the treatment of seed (fruits, tubers, grains) and nursery plants for the protection against fungal infections and against soil-borne phytopathogenic fungi.

Compounds of the formula I are active for example against phytopathogenic fungi which belong to the following classes: Fungi imperfecti (in particular Botrytis, furthermore Pyricularia, Helminthosporium, Fusarium, Septoria, Cercospora, Cercosporella and Alternaria); Basidiomycetes (for example Rhizoctonia, Hemileia, Puccinia); Ascomycetes (for example Venturia and Erysiphe, Podosphaera, Monilinia, Uncinula); Oomycetes (for example Phytophthora, Peronospora, Bremia, Pythium, Plasmopara).

Compounds of the formula I are furthermore valuable active ingredients against insects and Acarina which are found on useful plants and ornamentals in agriculture and horticulture, in particular in crops of rice, cotton, vegetables and fruit, and in forests, and are well tolerated by warm-blooded species, fish and plants. The compounds I are particularly suitable for controlling insects in crops of rice, fruit and vegetables, in particular plant-injurious insects. Other fields of application for the active ingredients of the invention are the protection of stored products and materials and, in the hygiene sector, in particular the protection of domestic animals and productive livestock. The compounds of the formula I are active against all or individual development stages of normally sensitive, but also resistant, pest species. Their activity can become apparent, for example, from the destruction of the pests, either immediately or only after some time has elapsed, for example during molting, or from reduced oviposition and/or hatching rates.

The abovementioned animal pests include, for example:

from the order Lepidoptera, for example,

Acleris spp., Adoxophyes spp., Aegeria spp., Agrotis spp., Alabama argillaceae, Amylois spp., Anticarsia gemmatilis, Archips spp., Argyrotaenia spp., Autographa spp., Busseola fusca, Cadra cautella, Carposina nipponensis, Chilo spp., Choristoneura spp., Clysia

ambiguella, *Cnaphalocrocis* spp., *Cnephasia* spp., *Cochylis* spp., *Coleophora* spp., *Crocidolomia* binotalis, *Cryptophlebia* leucotreta, *Cydia* spp., *Diatraea* spp., *Diparopsis* castanea, *Earias* spp., *Ephestia* spp., *Eucosma* spp., *Eupoecilia* ambiguella, *Euproctis* spp., *Euxoa* spp., *Grapholita* spp., *Hedya* nubiferana, *Heliothis* spp., *Hellula* undalis, *Hyphantria* cunea, *Keiferia* lycopersicella, *Leucoptera* scitella, *Lithocollethis* spp., *Lobesia* botrana, *Lymantria* spp., *Lyonetia* spp., *Malacosoma* spp., *Mamestra* brassicae, *Manduca* sexta, *Operophtera* spp., *Ostrinia* nubilalis, *Pammene* spp., *Pandemis* spp., *Panolis* flammea, *Pectinophora* gossypiella, *Phthorimaea* operculella, *Pieris* rapae, *Pieris* spp., *Plutella* xylostella, *Prays* spp., *Scirpophaga* spp., *Sesamia* spp., *Sparganothis* spp., *Spodoptera* spp., *Synanthedon* spp., *Thaumetopoea* spp., *Tortrix* spp., *Trichoplusia* ni and *Yponomeuta* spp.;

from the order Coleoptera, for example,

*Agriotes* spp., *Anthonomus* spp., *Atomaria* linearis, *Chaetocnema* tibialis, *Cosmopolites* spp., *Curculio* spp., *Dermestes* spp., *Diabrotica* spp., *Epilachna* spp., *Eremnus* spp., *Leptinotarsa* decemlineata, *Lissorhoptrus* spp., *Melolontha* spp., *Oryzaephilus* spp., *Otiorynchus* spp., *Phlyctinus* spp., *Popillia* spp., *Psylliodes* spp., *Rhizopertha* spp., *Scarabeidae*, *Sitophilus* spp., *Sitotroga* spp., *Tenebrio* spp., *Tribolium* spp. and *Trogoderma* spp.;

from the order Orthoptera, for example,

*Blatta* spp., *Blattella* spp., *Gryllotalpa* spp., *Leucophaea* maderae, *Locusta* spp., *Periplaneta* spp. and *Schistocerca* spp.;

from the order Isoptera, for example,

*Reticulitermes* spp.;

from the order Psocoptera, for example,

*Liposcelis* spp.;

from the order Anoplura, for example,

*Haematopinus* spp., *Linognathus* spp., *Pediculus* spp., *Pemphigus* spp. and *Phylloxera* spp.;

from the order Mallophaga, for example,

*Damalinea* spp. and *Trichodectes* spp.;

from the order Thysanoptera, for example,

*Frankliniella* spp., *Hercinothrips* spp., *Taeniothrips* spp., *Thrips* palmi, *Thrips* tabaci and *Scirtothrips* aurantii;

from the order Heteroptera, for example,

*Cimex* spp., *Distantiella* theobroma, *Dysdercus* spp., *Euchistus* spp., *Eurygaster* spp., *Leptocoris* spp., *Nezara* spp., *Piesma* spp., *Rhodnius* spp., *Sahlbergella* singularis, *Scotinophara* spp. and *Triatoma* spp.;

- 20 -

from the order Homoptera, for example,

*Aleurothrixus floccosus*, *Aleyrodes brassicae*, *Aonidiella* spp., *Aphididae*, *Aphis* spp., *Aspidiotus* spp., *Bemisia tabaci*, *Ceroplaster* spp., *Chrysomphalus aonidium*, *Chrysomphalus dictyospermi*, *Coccus hesperidum*, *Empoasca* spp., *Eriosoma larigerum*, *Erythroneura* spp., *Gascardia* spp., *Laodelphax* spp., *Lecanium corni*, *Lepidosaphes* spp., *Macrosiphus* spp., *Myzus* spp., *Nephotettix* spp., *Nilaparvata* spp., *Paratoria* spp., *Pemphigus* spp., *Planococcus* spp., *Pseudaulacaspis* spp., *Pseudococcus* spp., *Psylla* spp., *Pulvinaria aethiopica*, *Quadraspidiotus* spp., *Rhopalosiphum* spp., *Saissetia* spp., *Scaphoideus* spp., *Schizaphis* spp., *Sitobion* spp., *Trialeurodes vaporariorum*, *Trioza erytrae* and *Unaspis citri*;

from the order Hymenoptera, for example,

*Acromyrmex*, *Atta* spp., *Cephus* spp., *Diprion* spp., *Diprionidae*, *Gilpinia polytoma*, *Hoplocampa* spp., *Lasius* spp., *Monomorium pharaonis*, *Neodiprion* spp., *Solenopsis* spp. and *Vespa* spp.;

from the order Diptera, for example,

*Aedes* spp., *Antherigona soccata*, *Bibio hortulanus*, *Calliphora erythrocephala*, *Ceratitis* spp., *Chrysomyia* spp., *Culex* spp., *Cuterebra* spp., *Dacus* spp., *Drosophila melanogaster*, *Fannia* spp., *Gastrophilus* spp., *Glossina* spp., *Hypoderma* spp., *Hyppobosca* spp., *Liriomyza* spp., *Lucilia* spp., *Melanagromyza* spp., *Musca* spp., *Oestrus* spp., *Orseolia* spp., *Oscinella frit*, *Pegomyia hyoscyami*, *Phorbia* spp., *Rhagoletis pomonella*, *Sciara* spp., *Stomoxys* spp., *Tabanus* spp., *Tannia* spp. and *Tipula* spp.;

from the order Siphonaptera, for example,

*Ceratophyllus* spp. and *Xenopsylla cheopis*;

from the order Thysanura, for example,

*Lepisma saccharina* and

from the order Acarina, for example,

*Acarus siro*, *Aceria sheldoni*, *Aculus schlechtendali*, *Amblyomma* spp., *Argas* spp., *Boophilus* spp., *Brevipalpus* spp., *Bryobia praetiosa*, *Calipitimerus* spp., *Chorioptes* spp., *Dermanyssus gallinae*, *Eotetranychus carpini*, *Eriophyes* spp., *Hyalomma* spp., *Ixodes* spp., *Olygonychus pratensis*, *Ornithodoros* spp., *Panonychus* spp., *Phyllocoptruta oleivora*, *Polyphagotarsonemus latus*, *Psoroptes* spp., *Rhipicephalus* spp., *Rhizoglyphus* spp., *Sarcoptes* spp., *Tarsonemus* spp. and *Tetranychus* spp..

Target crops for crop-protecting use are, within the scope of the invention, for example the following types of plants: cereals (wheat, barley, rye, oats, rice, maize, sorghum and related species); beet (sugar and fodder beet); pome fruit, stone fruit and soft fruit (apples, pears, plums, peaches, almonds, cherries, strawberries, raspberries and blackberries);

pulses (beans, lentils, peas, soya beans); oil crops (oilseed rape, mustard, poppy, olives, sunflowers, coconut, castor, cocoa, groundnuts); cucurbits (pumpkin, cucumbers, melons); fibre plants (cotton, flax, hemp, jute); citrus fruit (oranges, lemons, grapefruit, tangerines); vegetables (spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes, bell pepper); the laurel family (avocado, cinammon, camphor), and plants such as tobacco, nuts, coffee, egg plants, sugarcane, tea, pepper, vines, hops, the banana family, latex plants and ornamentals.

The action of the compounds of the formula I according to the invention and of the compositions comprising them can be broadened considerably and adapted to prevailing circumstances by adding other microbicides, insecticides and/or acaricides.

Representatives of the following active ingredient classes are examples of suitable additives: organophosphorus compounds, nitrophenols and derivatives, formamidines, ureas, carbamates, pyrethroids, chlorinated hydrocarbons and *Bacillus thuringiensis* preparations.

As a rule, active ingredients of the formula I are used in the form of compositions and can be applied to the area or plants to be treated simultaneously or in succession with other active ingredients. These other active ingredients can be fertilizers, trace element mediators or other preparations which have an effect on plant growth. Selective herbicides and also insecticides, fungicides, bactericides, nematocides, molluscicides or mixtures of a plurality of these preparations, with or without other carriers conventionally used in the art of formulation, surfactants or other application-enhancing additives, can also be used.

Suitable carriers and additives can be solid or liquid and are those substances which are expedient in the art of formulation, for example natural or regenerated mineral substances, solvents, dispersants, wetting agents, tackifiers, thickeners, binders or fertilizers.

The following are suitable solvents: aromatic hydrocarbons, preferably the fractions C<sub>8</sub> to C<sub>12</sub>, for example xylene mixtures or substituted naphthalenes, phthalates such as dibutyl phthalate or dioctyl phthalate, aliphatic hydrocarbons such as cyclohexane or paraffins, alcohols and glycols as well as their ethers and esters, such as ethanol, ethylene glycol, ethylene glycol monomethyl ether or ethyl ether, ketones such as cyclohexanone, strongly polar solvents, such as N-methyl-2-pyrrolidone, dimethyl sulfoxide or dimethylformamide, and unepoxidized or epoxidized vegetable oils, such as epoxidized coconut oil or soya oil; or water.

Solid carriers which are generally used, for example for dusts and dispersible powders, are ground natural rocks, such as calcite, talc, kaolin, montmorillonite or attapulgite.

Particularly advantageous application enhancing additives which may result in a greatly reduced rate of application are, moreover, natural (animal or vegetable) or synthetic phospholipids from the series of the cephalins and lecithins, which can be obtained from, for example, soya beans.

Suitable surface-active compounds are, depending on the nature of the active ingredient of the formula I to be formulated, non-ionic, cationic and/or anionic surfactants which have good emulsifying, dispersing and wetting properties. Surfactants are also to be understood as meaning surfactant mixtures.

Suitable anionic surfactants can be so-called water-soluble salts and also water-soluble synthetic surface-active compounds.

Soaps are the alkali metal salts, alkaline earth metal salts or substituted or unsubstituted ammonium salts of higher fatty acids ( $C_{10}$ - $C_{22}$ ), for example the sodium salts or potassium salts of oleic or stearic acid, or of natural mixtures of fatty acids which can be obtained from, for example, coconut oil or tallow oil. The fatty acid methyltaurates must also be mentioned.

Suitable non-ionic surfactants are polyglycol ether derivatives of aliphatic or cycloaliphatic alcohols, saturated or unsaturated fatty acids and alkylphenols which have 3 to 30 glycol ether groups and 8 to 20 carbon atoms in the (aliphatic) hydrocarbon radical and 6 to 18 carbon atoms in the alkyl radical of the alkylphenols.

Examples of non-ionic surfactants are nonylphenol polyethoxyethanols, castor oil polyglycol ethers, polypropylene/polyethylene oxide adducts, tributylphenoxy-polyethoxyethanol, polyethylene glycol and octylphenoxypolyethoxyethanol.

Other suitable substances are fatty acid esters of polyoxyethylene sorbitan, such as polyoxyethylene sorbitan trioleate.

The cationic surfactants are mainly quaternary ammonium salts which have, as N-substituents, at least one alkyl radical having 8 to 22 C atoms and, as further substituents, lower, halogenated or free alkyl, benzyl or lower hydroxyalkyl radicals.



As a rule, the agrochemical preparations comprise 0.1 to 99 %, in particular 0.1 to 95 %, of active ingredient of the formula I, 99.9 to 1 %, in particular 99.9 to 5 %, of a solid or liquid additive and 0 to 25 %, in particular 0.1 to 25 %, of a surfactant.

While concentrated compositions are more preferred as commercial products, the end consumer will, as a rule, use dilute compositions.

The compositions can also comprise other additives such as stabilizers, antifoams, viscosity regulators, binders, tackifiers, fertilizers or other active ingredients for achieving specific effects.

The formulations, i.e. the compositions, preparations or combinations comprising the active ingredient of the formula I together or without a solid or liquid additive, are produced in a known manner, for example by intimately mixing and/or grinding the active ingredient with an extender, for example a solvent (mixture), a solid carrier and, if appropriate, surface-active compounds (surfactants).

A preferred process for applying an active ingredient of the formula I or an agrochemical composition which comprises at least one of these active ingredients, is application to the foliage (foliar application). Frequency and rate of application depend on the danger of attack by the pathogen in question. However, the active ingredients of the formula I can also reach the plant via the soil through the root system (systemic action) by drenching the locus of the plant with a liquid preparation or by incorporating the substances in solid form into the soil, for example in the form of granules (soil application). In the case of paddy rice, such granules can be metered out into the flooded rice field. Alternatively, the compounds of the formula I can be applied to seed kernels (coating), either by soaking the kernels in a liquid preparation of the active ingredient or by coating them with a solid preparation. In principle, any type of plant propagation material can be protected using the compounds of the formula I, for example seed, roots or stalks.

The compounds of the formula I are used as pure compounds or, preferably, together with the auxiliaries conventionally used in the art of formulation. To this end, they are advantageously processed in the known manner, for example to give emulsion concentrates, spreadable pastes, directly sprayable or dilutable solutions, dilute emulsions, wettable powders, soluble powders, dusts, granules, and encapsulations, for example in polymeric substances. The methods of application, such as spraying, atomizing, dusting,

- 24 -


broadcasting, painting on or pouring, as well as the nature of the compositions are selected to suit the intended aims and the prevailing circumstances. Favourable rates of application are, as a rule, 1 g to 2 kg of active ingredient (a.i.) per ha, preferably 25 g to 800 g of a.i./ha, particularly preferably 50 g to 400 g of a.i./ha. When used as seed dressing products, dosage rates of 0.001 g to 1.0 g of active ingredient are advantageously used per kg of seed.

The examples which follow are intended to illustrate the invention in greater detail without imposing any restriction thereto.

### Preparation Examples

#### Example H-1:


Preparation of



1.32 g of hydroxylamine hydrochloride are added to a solution of 5.2 g of methyl 3-methoxy-2-(((3-oxo-2-butyl)imino)oxy)-o-tolylacrylate (EP-A-370 629, No. 156) in 20 ml of pyridine. After the mixture has been stirred for 6 hours at 30°C, ice-water is added, the crystal cake which has formed after a few hours is filtered off, and the crystals are washed with water. Recrystallization from ethanol/water gives 4.8 g of methyl 3-methoxy-2-(((3-hydroximino-2-butyl)imino)oxy)-o-tolylacrylate (Comp. No. 1.73) as yellow crystals of melting point 104-107°C.

#### Example H-2:

Preparation of




- 25 -

In a pressure pipe, 5 ml of dimethylformamide are added to 0.2 g of an approximately 65 % sodium hydride suspension washed with hexane. Then, 1.6 g of methyl 3-methoxy-2-[[[(3-hydroximino-2-butyl)imino]oxy]o-tolyl]acrylate and hereupon 1.1 g of 2,2,2-trifluoroethyl iodide are added. After the evolution of hydrogen has subsided, the pressure pipe is sealed, and the mixture is stirred for 5 hours at 50°C. Then, the reaction solution is poured into ice-water and extracted using ethyl acetate, and the product is chromatographed on silica gel using ethyl acetate/hexane (1:3). This gives 1.2 g of methyl 3-methoxy-2-[[[(3-(2,2,2-trifluoroethoxyimino)-2-butyl)imino]oxy]o-tolyl]acrylate (Comp. No. 1.74) as a colourless oil.

<sup>1</sup>H NMR in CDCl<sub>3</sub>: chemical shift of the two imino-substituted methyl groups: 1.99 and 2.04 ppm.

#### Example H-3:

Preparation of



A solution of 15.0 g of p-methylpropiophenone and 20 ml of isopentyl nitrite in 30 ml of methanol is slowly added dropwise to 20 ml of sodium methylate (30 % in methanol). After the mixture has been stirred for 5 hours, 30 ml of water are added dropwise, and the mixture is subsequently acidified using acetic acid. The mixture is extracted using ethyl acetate, and the extract is washed with water, dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated in vacuo. The residue is recrystallized from diethyl ether/n-hexane. This gives α-hydroxyimino-4-methylpropiophenone as colourless crystals of melting point 122-124°C.

8.1 g of the previously obtained compound and 4.6 g of o-methylhydroxylamine hydrochloride in 40 ml of pyridine are refluxed for 1 hour. After an addition of toluene, the mixture is concentrated in vacuo, treated with water and extracted using ethyl acetate. The organic phase is washed with water, dried over sodium sulfate and concentrated in vacuo. α-Hydroxyimino-4-methylpropiophenone O-methyl oxime are crystallized from


- 26 -

diethyl ether/n-hexane in the form of colourless crystals of melting point 156-157°C.

1.87 g of the previously obtained oxime and 2.58 g of methyl 2-( $\alpha$ -bromo-o-tolyl)-3-methoxyacrylate are added to a suspension of 0.41 g of sodium hydride in 25 ml of N,N-dimethylformamide, and the reaction mixture is stirred for 3 hours. It is then acidified using acetic acid, treated with water and extracted using ethyl acetate. The organic phase is washed twice using saturated sodium hydrogen carbonate solution and once using saturated sodium chloride solution. After drying over anhydrous sodium sulfate, the solvent is distilled off in vacuo. H-3 crystallizes from diethyl ether/n-hexane in the form of colorless crystals of melting point 98-101°C (Comp. No. 1.117).

Example H-4:

Preparation of




0.37 g of an approximately 65 % sodium hydride suspension is washed with hexane, and 10 ml of dimethylformamide are added. A mixture of 2.59 g of methyl 2-(2-bromomethylphenyl)glyoxylate O-methyloxime and 2.42 g of 2-(diphenylhydrazono)-3-hydroximinobutane is added to this suspension, and the mixture is subsequently heated to 40-50°C using a hot-air blower until hydrogen is evolved vigorously. The mixture is then stirred for one hour with the exclusion of moisture, and poured into ice-water. Extraction using ethyl acetate and chromatography on silica gel using ethyl acetate/hexane (1:2) gives 3.7 g of methyl 2-(((3-diphenylhydrazono-2-butyl)imino)oxy)-o-tolylglyoxylate O-methyloxime (Comp. No. 2.19) as a yellow oil.  $^1\text{H}$  NMR in  $\text{CDCl}_3$ : chemical shift of the two imino-substituted methyl groups: 1.64 and 2.12 ppm.

- 27 -

Example H-5:


Preparation of



1.9 g of methyl 2-[[[(3-diphenylhydrazono-2-butyl)-imino]oxy]o-tolyl]glyoxylate O-methyloxime are stirred for 2 hours at room temperature in 10 ml of a 33 % methylamine solution in ethanol. Ethanol and the excess of methylamine are distilled off. The residue is taken up in diethyl ether, the solution is filtered, the filtrate is evaporated to dryness, the product remaining as a crystalline solid. Washing with hexane gives 1.8 g of N-monomethyl 2-[[[(3-diphenylhydrazono-2-butyl)imino]oxy]o-tolyl]glyoxylamide O-methyloxime (Comp. No. 3.19) as pale beige crystals of melting point 135-136°C.

Example H-6:


Preparation of




1.74 g of 1-[4-(2,2-dichlorocyclopropylmethoxy)phenyl]propane-1,2-dione 1-(O-methyloxime)-2-oxime, of the formula (AA), are added to 0.13 g of sodium hydride in 25 ml of N,N-dimethylformamide. 1.5 g of methyl 2-(2-bromomethylphenyl)-3-methoxyacrylate, of the formula (BB), are subsequently added, and the mixture is stirred for 3 hours at room temperature. The mixture is acidified using acetic acid, and water and ethyl acetate are added. The aqueous phase is separated off, and the organic phase is washed once using saturated sodium hydrogen carbonate solution and dried over anhydrous sodium sulfate. The solvent is evaporated in vacuo, and the residue obtained is purified on silica gel using hexane/ethyl acetate (3:1). The title compound is obtained as an oil. Separation by column chromatography on silica gel (hexane:ethyl acetate solution 3:1)

- 28 -

gives three isomers: isomer A, m.p. 86-88°C, isomer B, oil, isomer C, oil (Compound 5.9).



(AA)




(BB)

The following compounds can be prepared in this manner or analogously to one of the methods described further above:

(Abbreviations: Me=methyl, Et=ethyl,  $\Delta$ =cyclopropyl, Ph=phenyl, m.p.=melting point).

NMR: The chemical shifts are given as  $\delta$  (ppm) in  $\text{CDCl}_3$ .

Table 1



Ex. No.	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
1.1	NMe	Me	Me	6-CF <sub>3</sub> -2-pyridyl	2.17/2.19
1.2	NMe	H	Me	6-CF <sub>3</sub> -2-pyridyl	
1.3	NMe	Me	Δ	6-CF <sub>3</sub> -2-pyridyl	
1.4	NMe	Me	H	phenyl	2.09/2.18
1.5	NMe	Me	Me	phenyl	
1.6	NMe	Δ	Me	phenyl	
1.7	NMe	Me	Me	4-CF <sub>3</sub> -2-pyridyl	m.p. 100°C
1.8	NMe	H	Me	4-CF <sub>3</sub> -2-pyridyl	
1.9	NMe	Δ	Δ	phenyl	
1.10	NMe	Me	Me	5-CF <sub>3</sub> -2-pyridyl	2.17/2.18
1.11	NMe	H	Me	5-CF <sub>3</sub> -2-pyridyl	
1.12	-	H	Me	4-(1,2,4-triazolyl)	
1.13	-	Me	Me	4-(1,2,4-triazolyl)	161-163°C
1.14	-	Me	Δ	4-(1,2,4-triazolyl)	
1.15	-	Me	Δ	4-morpholinyl	
1.16	-	Me	Me	4-morpholinyl	113°C
1.17	-	H	Me	4-morpholinyl	
1.18	NPh*	H	Me	phenyl	
1.19	NPh*	Me	Me	phenyl	1.74/2.22
1.20	NPh*	Me	Δ	phenyl	
1.21	NPh*	Δ	Me	phenyl	
1.22	NMe	Me	Me	2-nitrophenyl	1.93/2.06
1.23	NMe	H	Me	2-nitrophenyl	
1.24	O	Me	Me	Me	76-78°C
1.25	O	H	Me	Me	1.97/7.68

- 30 -

Ex. No.	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
1.26	O	Δ	Me	Me	1.96/2.00
1.27	O	Me	Δ	Me	60-62°C
1.28	O	Me	H	Me	
1.29	NMe	H	Me	3-CF <sub>3</sub> -2-pyridyl	
1.30	NMe	Me	Me	3-CF <sub>3</sub> -2-pyridyl	2.06/2.18
1.31	NMe	Δ	Me	3-CF <sub>3</sub> -2-pyridyl	
1.32	NMe	Δ	Me	3-nitro-2-pyridyl	
1.33	NMe	H	Me	3-nitro-2-pyridyl	
1.34	NMe	Me	Me	3-nitro- 2-pyridyl	1.94/2.26
1.35	NMe	Me	Me	2-CF <sub>3</sub> -phenyl	1.85/2.10
1.36	NMe	Me	Δ	2-CF <sub>3</sub> -phenyl	
1.37	NMe	H	Me	2-CF <sub>3</sub> -phenyl	
1.38	NMe	Me	Me	3-CF <sub>3</sub> -phenyl	
1.39	NMe	Me	Δ	3-CF <sub>3</sub> -phenyl	
1.40	NMe	H	Me	4-CF <sub>3</sub> -phenyl	
1.41	NMe	Me	Me	4-CF <sub>3</sub> -phenyl	
1.42	NMe	Me	Me	2-chlorophenyl	1.73/2.13
1.43	NMe	Me	Me	3-chlorophenyl	2.12/2.17
1.44	NMe	H	Me	4-chlorophenyl	
1.45	NMe	Me	Me	4-chlorophenyl	2.10/2.15
1.46	O	Me	Me	phenyl	2.14/2.22
1.47	O	Me	Δ	phenyl	
1.48	O	Me	Me	benzyl	106-108°C
1.49	O	Me	Me	Et	1.92/1.95
1.50	O	H	Me	Et	
1.51	O	Δ	Me	Et	
1.52	O	Me	Δ	Et	
1.53	O	Me	H	Et	
1.54	O	H	Me	methoxymethyl	
1.55	O	Me	Me	methoxymethyl	83-84°C
1.56	O	Me	Δ	methoxymethyl	



- 31 -

Ex. No.	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
1.57	O	Δ	Me	methoxymethyl	
1.58	O	Me	Me	ethoxymethyl	2.04/2.04
1.59	O	H	Me	cyanomethyl	
1.60	O	Me	Me	cyanomethyl	2.04/2.04
1.61	O	Δ	Me	cyanomethyl	
1.62	-	Me	Me	azepino	2.02/2.04
1.63	-	Me	Me	piperidino	
1.64	-	Me	Me	pyrrolidino	
1.65	O	H	Me	tert-butyl	
1.66	O	Me	Me	tert-butyl	1.96/2.04
1.67	O	Me	Me	propargyl	114-115°C
1.68	O	Δ	Me	propargyl	
1.69	O	Me	Δ	propargyl	
1.70	O	Me	Me	2,2-dichloro- cyclopropylmethyl	2.02/2.02
1.71	O	Δ	Me	2,2-dichloro- cyclopropylmethyl	
1.72	O	H	Me	H	
1.73	O	Me	Me	H	104-107°C
1.74	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub>	1.99/2.04
1.75	O	Δ	Me	CF <sub>3</sub> CH <sub>2</sub>	
1.76	O	Me	H	CF <sub>3</sub> CH <sub>2</sub>	
1.77	O	Me	H	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	
1.78	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	
1.79	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	2.10/2.13
1.80	O	Δ	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	
1.81	NMe	Me	Me	Me	94-98°C
1.82	NMe	Me	Δ	Me	
1.83	O	Me	Me	CH <sub>2</sub> -CCl=CH <sub>2</sub>	80°C
1.84	O	Δ	Me	CH <sub>2</sub> -CCl=CH <sub>2</sub>	
1.85	O	Me	Me	propyl	
1.86	O	Me	Me	butyl	36-38°C

- 32 -

Ex. No.	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
1.87	O	Me	Me	hexyl	1.99/2.01
1.88	O	Me	Me	methoxycarbonyl- methyl	90-93°C
1.89	O	H	Me	methoxycarbonyl- methyl	
1.90	O	Me	Me	3-fluorobenzyl	2.00/2.04
1.91	O	Me	Me	4-chlorobenzyl	
1.92	O	Me	Me	2-chlorobenzyl	
1.93	O	Me	Me	2-CF <sub>3</sub> -benzyl	
1.94	O	Me	Me	3-CF <sub>3</sub> -benzyl	56-58°C
1.95	O	Me	Me	4-CF <sub>3</sub> -benzyl	
1.96	O	Me	Me	3,4-dichlorobenzyl	
1.97	O	Me	Me	2,4,6-trimethylbenzyl	
1.98	O	Me	Me	4-chloro-2-nitrobenzyl	
1.99	O	Me	Me	3-methoxybenzyl	
1.100	O	Me	Me	2-phenethyl	1.97/2.02
1.101	O	Me	Me	3-phenylpropyl	
1.102	O	Me	Me	2-(4-nitrophenyl)ethyl	
1.103	O	Me	Me	2-(2-CF <sub>3</sub> -phenyl)ethyl	
1.104	O	Me	Me	2-(4-methoxyphenyl)ethyl	
1.105	O	Me	Me	2-chloro-6-fluorobenzyl	
1.106	O	Me	Me	3,4-methylenedioxybenzyl	
1.107	O	Me	Me	2-cyanobenzyl	
1.108	O	Me	Me	2-(4-chlorophenyl)ethyl	
1.109	O	Me	Me	cyclopropyl- methyl	2.01/2.01
1.110	O	Me	Me	2-(1,3-dioxolanyl)methyl	
1.111	O	Me	Me	2,2,3,3-tetrafluorocyclo- butylmethyl	
1.112	O	Me	Me	α-fluoroethoxy carbonylmethyl	
1.113	O	Me	2-thienyl	Me	2.06/2.10/


- 33 -

Ex. No.	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
1.114	O	Me	4-methyl-phenyl	Et	2.20 (E/Z) 2.12
1.115	NMe	Me	4-methyl-phenyl	Me	2.10 (isomer1) 2.08 (isomer2)
1.116	O	Me	Me	CH <sub>2</sub> FCH <sub>2</sub>	50-51°C
1.117	O	Me	Me	2-(4-morpho- lino)ethyl	1.99/2.01
1.118	-	Me	Me	1-pyrrol	89-91°C
1.119	O	Me	Me	2-(1-piperidino)- ethyl	
1.120	O	Me	Me	2-fluorobenzyl	93-96°C
1.121	O	Me	Me	4-fluorobenzyl	104-106°C

\*\* m.p. or <sup>1</sup>H NMR of R<sub>1</sub>/R<sub>2</sub> or R<sub>3</sub>

- 34 -

Table 2



Ex. No.	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
2.1	NMe	Me	Me	6-CF <sub>3</sub> -2-pyridyl	87-88°C
2.2	NMe	H	Me	6-CF <sub>3</sub> -2-pyridyl	
2.3	NMe	Me	Δ	6-CF <sub>3</sub> -2-pyridyl	
2.4	NMe	Me	H	phenyl	
2.5	NMe	Me	Me	phenyl	
2.6	NMe	Δ	Me	phenyl	
2.7	NMe	Me	Me	4-CF <sub>3</sub> -2-pyridyl	
2.8	NMe	H	Me	4-CF <sub>3</sub> -2-pyridyl	
2.9	NMe	Δ	Δ	phenyl	
2.10	NMe	Me	Me	5-CF <sub>3</sub> -2-pyridyl	
2.11	NMe	H	Me	5-CF <sub>3</sub> -2-pyridyl	1.64/2.12
2.12	-	H	Me	4-(1,2,4-triazolyl)	
2.13	-	Me	Me	4-(1,2,4-triazolyl)	
2.14	-	Me	Δ	4-(1,2,4-triazolyl)	
2.15	-	Me	Δ	4-morpholinyl	
2.16	-	Me	Me	4-morpholinyl	
2.17	-	H	Me	4-morpholinyl	
2.18	NPh	H	Me	phenyl	
2.19	NPh	Me	Me	phenyl	
2.20	NPh	Me	Δ	phenyl	
2.21	NPh	Δ	Me	phenyl	104-106°C
2.22	NMe	Me	Me	2-nitrophenyl	
2.23	NMe	H	Me	2-nitrophenyl	
2.24	O	Me	Me	Me	
2.25	O	H	Me	Me	

- 35 -

Ex. No.	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
2.26	O	Δ	Me	Me	1.92/1.97
2.27	O	Me	Δ	Me	67-70°C
2.28	O	Me	H	Me	
2.29	NMe	H	Me	3-CF <sub>3</sub> -2-pyridyl	
2.30	NMe	Me	Me	3-CF <sub>3</sub> -2-pyridyl	
2.31	NMe	Δ	Me	3-CF <sub>3</sub> -2-pyridyl	
2.32	NMe	Δ	Me	3-nitro-2-pyridyl	
2.33	NMe	H	Me	3-nitro-2-pyridyl	
2.34	NMe	Me	Me	3-nitro-2-pyridyl	
2.35	NMe	Me	Me	2-CF <sub>3</sub> -phenyl	
2.36	NMe	Me	Δ	2-CF <sub>3</sub> -phenyl	
2.37	NMe	H	Me	2-CF <sub>3</sub> -phenyl	
2.38	NMe	Me	Me	3-CF <sub>3</sub> -phenyl	
2.39	NMe	Me	Δ	3-CF <sub>3</sub> -phenyl	
2.40	NMe	H	Me	4-CF <sub>3</sub> -phenyl	
2.41	NMe	Me	Me	4-CF <sub>3</sub> -phenyl	
2.42	NMe	Me	Me	2-chlorophenyl	81°C
2.43	NMe	Me	Me	3-chlorophenyl	2.10/2.13
2.44	NMe	H	Me	4-chlorophenyl	
2.45	NMe	Me	Me	4-chlorophenyl	2.08/2.13
2.46	O	Me	Me	phenyl	89-90°C
2.47	O	Me	Δ	phenyl	
2.48	O	Me	Me	benzyl	
2.49	O	Me	Me	Et	
2.50	O	H	Me	Et	
2.51	O	Δ	Me	Et	
2.52	O	Me	Δ	Et	
2.53	O	Me	H	Et	
2.54	O	H	Me	methoxymethyl	
2.55	O	Me	Me	methoxymethyl	87-88°C
2.56	O	Me	Δ	methoxymethyl	
2.57	O	Δ	Me	methoxymethyl	

- 36 -

Ex. No.	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
2.58	O	Me	Me	ethoxymethyl	
2.59	O	H	Me	cyanomethyl	
2.60	O	Me	Me	cyanomethyl	94-95°C
2.61	O	Δ	Me	cyanomethyl	
2.62	-	Me	Me	azepino	
2.63	-	Me	Me	piperidino	
2.64	-	Me	Me	pyrrolidino	
2.65	O	H	Me	tert-butyl	
2.66	O	Me	Me	tert-butyl	1.93/2.02
2.67	O	Me	Me	propargyl	120-122°C
2.68	O	Δ	Me	propargyl	
2.69	O	Me	Δ	propargyl	
2.70	O	Me	Me	2,2-dichloro- cyclo- propylmethyl	1.99/1.99
2.71	O	Δ	Me	2,2-dichlorocyclo- propylmethyl	
2.72	O	H	Me	H	
2.73	O	Me	Me	H	114-116°C
2.74	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub>	52-53°C
2.75	O	Δ	Me	CF <sub>3</sub> CH <sub>2</sub>	
2.76	O	Me	H	CF <sub>3</sub> CH <sub>2</sub>	
2.77	O	Me	H	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	
2.78	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	
2.79	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	86°C
2.80	O	Δ	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	
2.81	NMe	Me	Me	Me	80-81°C
2.82	NMe	Me	Δ	Me	
2.83	O	Me	Me	CH <sub>2</sub> -CCl=CH <sub>2</sub>	66°C
2.84	O	Δ	Me	CH <sub>2</sub> -CCl=CH <sub>2</sub>	
2.85	O	Me	Me	propyl	
2.86	O	Me	Me	butyl	58-62°C

- 37 -

Ex. No.	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
2.87	O	Me	Me	hexyl	50-53°C
2.88	O	Me	Me	methoxycarbonyl- methyl	1.96/2.04
2.89	O	H	Me	methoxycarbonyl- methyl	
2.90	O	Me	Me	3-fluorobenzyl	
2.91	O	Me	Me	4-chlorobenzyl	
2.92	O	Me	Me	2-chlorobenzyl	
2.93	O	Me	Me	2-CF <sub>3</sub> -benzyl	
2.94	O	Me	Me	3-CF <sub>3</sub> -benzyl	87-88°C
2.95	O	Me	Me	4-CF <sub>3</sub> -benzyl	
2.96	O	Me	Me	3,4-dichlorobenzyl	
2.97	O	Me	Me	2,4,6-trimethylbenzyl	
2.98	O	Me	Me	4-chloro-2-nitrobenzyl	
2.99	O	Me	Me	3-methoxybenzyl	
2.100	O	Me	Me	2-phenethyl	
2.101	O	Me	Me	3-phenylpropyl	
2.102	O	Me	Me	2-(4-nitrophenyl)ethyl	
2.103	O	Me	Me	2-(2-CF <sub>3</sub> -phenyl)ethyl	
2.104	O	Me	Me	2-(4-methoxyphenyl)ethyl	
2.105	O	Me	Me	2-chloro-6-fluorobenzyl	
2.106	O	Me	Me	3,4-methylenedioxybenzyl	
2.107	O	Me	Me	2-cyanobenzyl	
2.108	O	Me	Me	2-(4-chlorophenyl)ethyl	
2.109	O	Me	Me	cyclopropylmethyl	1.98/1.98
2.110	O	Me	Me	2-(1,3-dioxolanyl)methyl	
2.111	O	Me	Me	2,2,3,3-tetrafluorocyclo- butylmethyl	
2.112	O	Me	Me	α-fluoroethoxycarbonylmethyl	
2.113	O	Me	2-thienyl	Me	83-86°C
2.114	O	Me	4-methyl- phenyl	Et	94-96°C

- 38 -


Ex. No.	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
2.115	NMe	Me	4-methyl- phenyl	Me	
2.116	O	Me	Me	CH <sub>2</sub> FCH <sub>2</sub>	62-63°C
2.117	O	Me	Me	2-(4-morpho- lino)ethyl	1.97/1.99
2.118	-	Me	Me	1-pyrrol	
2.119	O	Me	Me	2-(1-piperidino)- ethyl	1.95/2.00
2.120	O	Me	Me	2-fluorobenzyl	81-83°C
2.121	O	Me	Me	4-fluorobenzyl	116-118°C

\*\* m.p. or <sup>1</sup>H NMR of R<sub>1</sub>/R<sub>2</sub> or R<sub>3</sub>



- 39 -

Table 3



Ex. No.	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
3.1	NMe	Me	Me	6-CF <sub>3</sub> -2-pyridyl	2.07/2.15
3.2	NMe	H	Me	6-CF <sub>3</sub> -2-pyridyl	
3.3	NMe	Me	Δ	6-CF <sub>3</sub> -2-pyridyl	
3.4	NMe	Me	H	phenyl	
3.5	NMe	Me	Me	phenyl	
3.6	NMe	Δ	Me	phenyl	
3.7	NMe	Me	Me	4-CF <sub>3</sub> -2-pyridyl	
3.8	NMe	H	Me	4-CF <sub>3</sub> -2-pyridyl	
3.9	NMe	Δ	Δ	phenyl	
3.10	NMe	Me	Me	5-CF <sub>3</sub> -2-pyridyl	
3.11	NMe	H	Me	5-CF <sub>3</sub> -2-pyridyl	
3.12	-	H	Me	4-(1,2,4-triazolyl)	
3.13	-	Me	Me	4-(1,2,4-triazolyl)	
3.14	-	Me	Δ	4-(1,2,4-triazolyl)	
3.15	-	Me	Δ	4-morpholinyl	
3.16	-	Me	Me	4-morpholinyl	135-136°C
3.17	-	H	Me	4-morpholinyl	
3.18	NPh	H	Me	phenyl	
3.19	NPh	Me	Me	phenyl	
3.20	NPh	Me	Δ	phenyl	
3.21	NPh	Δ	Me	phenyl	
3.22	NMe	Me	Me	2-nitrophenyl	
3.23	NMe	H	Me	2-nitrophenyl	
3.24	O	Me	Me	Me	90-92°C
3.25	O	H	Me	Me	1.97/7.68

- 40 -

Ex. No.	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
3.26	O	Δ	Me	Me	1.94/1.95
3.27	O	Me	Δ	Me	88-90°C
3.28	O	Me	H	Me	
3.29	NMe	H	Me	3-CF <sub>3</sub> -2-pyridyl	
3.30	NMe	Me	Me	3-CF <sub>3</sub> -2-pyridyl	
3.31	NMe	Δ	Me	3-CF <sub>3</sub> -2-pyridyl	
3.32	NMe	Δ	Me	3-nitro-2-pyridyl	
3.33	NMe	H	Me	3-nitro-2-pyridyl	
3.34	NMe	Me	Me	3-nitro-2-pyridyl	
3.35	NMe	Me	Me	2-CF <sub>3</sub> -phenyl	
3.36	NMe	Me	Δ	2-CF <sub>3</sub> -phenyl	
3.37	NMe	H	Me	2-CF <sub>3</sub> -phenyl	
3.38	NMe	Me	Me	3-CF <sub>3</sub> -phenyl	
3.39	NMe	Me	Δ	3-CF <sub>3</sub> -phenyl	
3.40	NMe	H	Me	4-CF <sub>3</sub> -phenyl	
3.41	NMe	Me	Me	4-CF <sub>3</sub> -phenyl	
3.42	NMe	Me	Me	2-chlorophenyl	1.72/2.11
3.43	NMe	Me	Me	3-chlorophenyl	2.09/2.14
3.44	NMe	H	Me	4-chlorophenyl	
3.45	NMe	Me	Me	4-chlorophenyl	2.08/2.13
3.46	O	Me	Me	phenyl	
3.47	O	Me	Δ	phenyl	
3.48	O	Me	Me	benzyl	
3.49	O	Me	Me	Et	
3.50	O	H	Me	Et	
3.51	O	Δ	Me	Et	
3.52	O	Me	Δ	Et	
3.53	O	Me	H	Et	
3.54	O	H	Me	methoxymethyl	
3.55	O	Me	Me	methoxymethyl	2.00/2.03
3.56	O	Me	Δ	methoxymethyl	
3.57	O	Δ	Me	methoxymethyl	

- 41 -

Ex. No.	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
3.58	O	Me	Me	ethoxymethyl	
3.59	O	H	Me	cyanomethyl	
3.60	O	Me	Me	cyanomethyl	2.00/2.01
3.61	O	Δ	Me	cyanomethyl	
3.62	-	Me	Me	azepino	
3.63	-	Me	Me	piperidino	
3.64	-	Me	Me	pyrrolidino	
3.65	O	H	Me	tert-butyl	
3.66	O	Me	Me	tert-butyl	1.95/2.01
3.67	O	Me	Me	propargyl	114-115°C
3.68	O	Δ	Me	propargyl	
3.69	O	Me	Δ	propargyl	
3.70	O	Me	Me	2,2-dichloro- cyclopropyl- methyl	
3.71	O	Δ	Me	2,2-dichloro- cyclopropyl- methyl	
3.72	O	H	Me	H	
3.73	O	Me	Me	H	
3.74	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub>	94-96°C
3.75	O	Δ	Me	CF <sub>3</sub> CH <sub>2</sub>	
3.76	O	Me	H	CF <sub>3</sub> CH <sub>2</sub>	
3.77	O	Me	H	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	
3.78	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	
3.79	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	
3.80	O	Δ	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	
3.81	NMe	Me	Me	Me	90-91°C
3.82	NMe	Me	Δ	Me	
3.83	O	Me	Me	CH <sub>2</sub> -CCl=CH <sub>2</sub>	1.97/2.02
3.84	O	Δ	Me	CH <sub>2</sub> -CCl=CH <sub>2</sub>	
3.85	O	Me	Me	propyl	

- 42 -


Ex. No.	A	R <sub>1</sub> R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
3.86	O	Me Me	butyl	91-92°C
3.87	O	Me Me	hexyl	84-86°C
3.88	O	Me Me	methylcarbamoyl- methyl	127-129°C
3.89	O	H Me	methoxycarbonyl- methyl	
3.90	O	Me Me	3-fluorobenzyl	
3.91	O	Me Me	4-chlorobenzyl	
3.92	O	Me Me	2-chlorobenzyl	
3.93	O	Me Me	2-CF <sub>3</sub> -benzyl	
3.94	O	Me Me	3-CF <sub>3</sub> -benzyl	1.96/2.00
3.95	O	Me Me	4-CF <sub>3</sub> -benzyl	
3.96	O	Me Me	3,4-dichlorobenzyl	
3.97	O	Me Me	2,4,6-trimethylbenzyl	
3.98	O	Me Me	4-chloro-2-nitrobenzyl	
3.99	O	Me Me	3-methoxybenzyl	
3.100	O	Me Me	2-phenethyl	
3.101	O	Me Me	3-phenylpropyl	
3.102	O	Me Me	2-(4-nitrophenyl)ethyl	
3.103	O	Me Me	2-(2-CF <sub>3</sub> -phenyl)ethyl	
3.104	O	Me Me	2-(4-methoxyphenyl)ethyl	
3.105	O	Me Me	2-chloro-6-fluorobenzyl	
3.106	O	Me Me	3,4-methylenedioxybenzyl	
3.107	O	Me Me	2-cyanobenzyl	
3.108	O	Me Me	2-(4-chlorophenyl)ethyl	
3.109	O	Me Me	cyclopropylmethyl	2.00/2.00
3.110	O	Me Me	2-(1,3-dioxolanyl)methyl	
3.111	O	Me Me	2,2,3,3-tetrafluorocyclo- butylmethyl	
3.112	O	Me Me	α-fluoroethoxy- carbonylmethyl	
3.113	O	Me 2-thienyl	Me	

- 43 -

Ex. No.	A	R <sub>1</sub> R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical data**
3.114	O	Me 4-methylphenyl	Et	103-104°C
3.115	NMe	Me 4-methylphenyl	Me	
3.116	O	Me Me	CH <sub>2</sub> FCH <sub>2</sub>	95-97°C
3.117	O	Me Me	2-(4-morpho- lino)ethyl	1.98/2.06
3.118	-	Me Me	1-pyrrol	
3.119	O	Me Me	2-(1-piperidino)- ethyl	1.96/1.98
3.120	O	Me Me	2-fluorobenzyl-	90-91°C
3.121	O	Me Me	4-fluorobenzyl	130-131°C

\*\* m.p. or <sup>1</sup>H NMR of R<sub>1</sub>/R<sub>2</sub> or R<sub>3</sub>

Table 4:



Ex.	X	Y	R <sub>1</sub>	Z*	n	B or D	Physical data**
4.1	CH	OMe	Me	-	1	3-CF <sub>3</sub>	133-135°C
4.2	CH	OMe	Me	-	1	4-chloro	140-142°C
4.3	CH	OMe	Me	-	1	3-chloro	175-177°C
4.4	CH	OMe	Me	-	1	2-fluoro	136-137°C
4.5	CH	OMe	Me	-	1	4-methyl	isom.1: 98-101°C isom.2: oil isom.3: oil
4.6	CH	OMe	Me	O	0	methyl-	104-106°C
4.7	CH	OMe	Me	-	1	4-bromo	126-128°C
4.8	CH	OMe	Me	-	1	4-fluoro	141-144°C
4.9	CH	OMe	Me	-	2	3-F-5-CF <sub>3</sub>	164-166°C
4.10	CH	OMe	Me	-	0	-	141-143°C
4.11	CH	OMe	Me	-	1	2-methyl	113-115°C
4.12	CH	OMe	Me	-	1	3-bromo	176-178°C
4.13	CH	OMe	Me	-	2	3,4-methylenedioxy	143-145°C
4.14	CH	OMe	Me	O	0	allyl-	77-80°C (isomer1) 2.08(isomer2)
4.15	CH	OMe	SMe	-	1	4-methyl	2.08 (isomer1) 2.11 (isomer2)
4.16	CH	OMe	Et	-	1	4-methyl	3.90 (isomer1) 3.97 (isomer2)
4.17	CH	OMe	Me	-	1	4-isobutyl-	2.11 (isomer1) 2.08 (isomer2)
4.18	CH	OMe	Me	O	0	propargyl-	2.06/2.11(E/Z)

- 45 -

Ex.	X	Y	R <sub>1</sub>	Z*	n	B or D	Physical data**
4.19	CH	OMe	Me	O	0	2,2,2-tri-fluoroethyl	2.11 (isomer1) 2.07 (isomer2)
4.20	CH	OMe	Me	O	0	ethyl	93-95°C (isomer1) 2.06 (isomer2)
4.21	CH	OMe	CN	-	1	4-methyl	98-133°C(E/Z)
4.22	CH	OMe	CN	-	1	4-chloro	106-108°C
4.23	CH	OMe	CN	-	2	3,4-dichloro	128-130°C 109-111°C(E/Z)
4.24	CH	OMe	CN	O	0	-CF <sub>3</sub>	oil (E/Z)
4.25	CH	OMe	CN	-	1	3-CF <sub>3</sub>	
4.26	CH	OMe	CN	-	1	2-chloro	
4.27	CH	OMe	CN	-	1	4-fluoro	
4.28	CH	OMe	Me	O	0	phenyl	112-114°C(iso.1) oil (iso.2)
4.29	CH	OMe	Me	O	0	-CH <sub>2</sub> CH=CCl <sub>2</sub>	
4.30	CH	OMe	Me	O	0	CH <sub>2</sub> -CH=CF <sub>2</sub>	
4.31	CH	OMe	Me	O	0	-CH <sub>2</sub> CH=CBr <sub>2</sub>	
4.32	CH	OMe	Me	O	0	4-Cl-phenyl	
4.33	CH	OMe	Me	O	0	4-F-phenyl	
4.34	CH	OMe	Me	S	0	phenyl	
4.35	CH	OMe	Me	-CH <sub>2</sub> O-	0	phenyl ***	isomer 1: resin isomer 2: resin
4.36	CH	OMe	Me	-OCH <sub>2</sub> -	0	phenyl ***	oil
4.37	N	OMe	Me	-	1	3-CF <sub>3</sub>	
4.38	N	OMe	Me	-	1	4-chloro	
4.39	N	OMe	Me	-	1	3-chloro	
4.40	N	OMe	Me	-	1	2-fluoro	111-112°C
4.41	N	OMe	Me	-	1	4-methyl	101-103°C
4.42	N	OMe	Me	O	0	4-methyl-	104-106°C
4.43	N	OMe	Me	-	1	4-bromo	134-136°C
4.44	N	OMe	Me	-	1	4-fluoro	
4.45	N	OMe	Me	-	2	3-F-5-CF <sub>3</sub>	
4.46	N	OMe	Me	-	0	-	
4.47	N	OMe	Me	-	1	2-methyl	105-107°C

- 46 -

Ex.	X	Y	R <sub>1</sub>	Z*	n	B or D	Physical data**
4.48	N	OMe	Me	-	1	3-bromo	
4.49	N	OMe	Me	-	2	3,4-methylenedioxy	
4.50	N	OMe	Me	O	0	allyl	74-76°C
4.51	N	OMe	SMe	-	1	4-methyl	88-91°C
4.52	N	OMe	Et	-	1	4-methyl	
4.53	N	OMe	Me	-	1	4-isobutyl-	
4.54	N	OMe	Me	O	0	propargyl	oil
4.55	N	OMe	Me	O	0	2,2,2-trifluoroethyl	
4.56	N	OMe	Me	O	0	ethyl	oil
4.57	N	OMe	CN	-	1	4-methyl	
4.58	N	OMe	CN	-	1	4-chloro	
4.59	N	OMe	CN	-	2	3,4-dichloro	
4.60	N	OMe	CN	O	0	-CF <sub>3</sub>	
4.61	N	OMe	CN	-	1	3-CF <sub>3</sub>	
4.62	N	OMe	CN	-	1	2-chloro	
4.63	N	OMe	CN	-	1	4-fluoro	
4.64	N	OMe	Me	O	0	phenyl	
4.65	N	OMe	Me	O	0	-CH <sub>2</sub> CH=CCl <sub>2</sub>	
4.66	N	OMe	Me	O	0	-CH <sub>2</sub> CH=CF <sub>2</sub>	
4.67	N	OMe	Me	O	0	-CH <sub>2</sub> CH=CBr <sub>2</sub>	
4.68	N	OMe	Me	O	0	4-Cl-phenyl	
4.69	N	OMe	Me	O	0	4-F-phenyl	
4.70	N	OMe	Me	S	0	phenyl	
4.71	N	OMe	Me	-CH <sub>2</sub> O-	0	phenyl ***	
4.72	N	OMe	Me	-OCH <sub>2</sub> -	0	phenyl ***	
4.73	N	NHCH <sub>3</sub>	Me	-	1	3-CF <sub>3</sub>	
4.74	N	NHCH <sub>3</sub>	Me	-	1	4-chloro	
4.75	N	NHCH <sub>3</sub>	Me	-	1	3-chloro	
4.76	N	NHCH <sub>3</sub>	Me	-	1	2-fluoro	2.11
4.77	N	NHCH <sub>3</sub>	Me	-	1	4-methyl	2.10
4.78	N	NHCH <sub>3</sub>	Me	O	0	methyl-	
4.79	N	NHCH <sub>3</sub>	Me	-	1	4-bromo	2.09
4.80	N	NHCH <sub>3</sub>	Me	-	1	4-fluoro	
4.81	N	NHCH <sub>3</sub>	Me	-	2	3-fluoro-5-CF <sub>3</sub>	



- 47 -

Ex.	X	Y	R <sub>1</sub>	Z*	n	B or D	Physical data**
4.82	N	NHCH <sub>3</sub>	Me	-	0	-	
4.83	N	NHCH <sub>3</sub>	Me	-	1	2-methyl	2.05
4.84	N	NHCH <sub>3</sub>	Me	-	1	3-bromo	
4.85	N	NHCH <sub>3</sub>	Me	-	2	3,4-methylenedioxy	
4.86	N	NHCH <sub>3</sub>	Me	O	0	allyl	118-120°C
4.87	N	NHCH <sub>3</sub>	SMe	-	1	4-methyl	2.06
4.88	N	NHCH <sub>3</sub>	Et	-	1	4-methyl	
4.89	N	NHCH <sub>3</sub>	Me	-	1	4-isobutyl-	
4.90	N	NHCH <sub>3</sub>	Me	O	0	propargyl	oil
4.91	N	NHCH <sub>3</sub>	Me	O	0	2,2,2-trifluoroethyl	
4.92	N	NHCH <sub>3</sub>	Me	O	0	ethyl	oil
4.93	N	NHCH <sub>3</sub>	CN	-	1	4-methyl	
4.94	N	NHCH <sub>3</sub>	CN	-	1	4-chloro	
4.95	N	NHCH <sub>3</sub>	CN	-	2	3,4-dichloro	
4.96	N	NHCH <sub>3</sub>	CN	O	0	-CF <sub>3</sub>	
4.97	N	NHCH <sub>3</sub>	CN	-	1	3-CF <sub>3</sub>	
4.98	N	NHCH <sub>3</sub>	CN	-	1	2-chloro	
4.99	N	NHCH <sub>3</sub>	CN	-	1	4-fluoro	
4.100	N	NHCH <sub>3</sub>	Me	O	0	phenyl	
4.101	N	NHCH <sub>3</sub>	Me	O	0	-CH <sub>2</sub> CH=CCl <sub>2</sub>	
4.102	N	NHCH <sub>3</sub>	Me	O	0	-CH <sub>2</sub> CH=CF <sub>2</sub>	
4.103	N	NHCH <sub>3</sub>	Me	O	0	-CH <sub>2</sub> CH=CBr <sub>2</sub>	
4.104	N	NHCH <sub>3</sub>	Me	O	0	4-Cl-phenyl	
4.105	N	NHCH <sub>3</sub>	Me	O	0	4-F-phenyl	
4.106	N	NHCH <sub>3</sub>	Me	S	0	phenyl	
4.107	N	NHCH <sub>3</sub>	Me	-CH <sub>2</sub> O-	0	phenyl	***
4.108	N	NHCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	phenyl	***
4.109	CH	OCH <sub>3</sub>	Me	O	0	n-propyl	isomer 1: oil isomer 2: oil
4.110	CH	OCH <sub>3</sub>	Me	O	0	3,3-dimethyl allyl	107-110°C(iso 1) oil (iso 2)
4.111	CH	OCH <sub>3</sub>	Me	O	0	2-methyl- allyl	isomer 1: 97-99°C isomer 2: oil

- 48 -

Ex.	X	Y	R <sub>1</sub>	Z*	n	B or D	Physical data**
4.112	CH	OCH <sub>3</sub>	Me	O	0	3-methyl-allyl	isomer 1: oil isomer 2: oil
4.113	CH	OCH <sub>3</sub>	Me	O	0	allyl	isomer 1: 74-76°C
4.114	CH	OCH <sub>3</sub>	Me	O	0	propargyl	oil
4.115	CH	NHCH <sub>3</sub>	Me	O	0	allyl	
4.116	CH	NHCH <sub>3</sub>	Me	O	0	propargyl	
4.117	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	3-CF <sub>3</sub> -phenyl***	isomer 1: 83-85°C isomer 2: resin
4.118	CH	OCH <sub>3</sub>	Me	O	0	3-CF <sub>3</sub> -phenyl	isomer 1: 127-128 isomer 2: oil
4.119	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	C <sub>6</sub> H <sub>11</sub> ***	isomer 1: oil isomer 2: oil
4.120	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	phenyl***	isomer 1: oil isomer 2: oil
4.121	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	4-F-phenyl***	isomer 1: oil isomer 2: oil
4.122	CH	OCH <sub>3</sub>	Me	S	0	CH <sub>3</sub>	isomer 1: 119-121°C isomer 2: resin
4.123	CH	OCH <sub>3</sub>	Me	-SO <sub>2</sub> -	0	CH <sub>3</sub>	isomer 1: foam isomer 2: resin
4.124	CH	OCH <sub>3</sub>	Me	-CH <sub>2</sub> O-	0	phenyl***	isomer 1: oil isomer 2: resin
4.125	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	4-CF <sub>3</sub> -phenyl***	
4.126	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	4-Br-phenyl***	
4.127	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	4-CH <sub>3</sub> -phenyl***	
4.128	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	4-OCH <sub>3</sub> -phenyl***	
4.129	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	2-CF <sub>3</sub> -phenyl***	
4.130	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	2-F-phenyl***	isomer 1: oil isomer 2: oil
4.131	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	2-Cl-phenyl***	
4.132	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	2-Br-phenyl***	
4.133	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	3-F-phenyl***	isomer 1: 111-112°C isomer 2: oil
4.134	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	3-Cl-phenyl***	isomer 1: oil

- 49 -

Ex.	X	Y	R <sub>1</sub>	Z*	n	B or D	Physical data**
4.135	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	3-Br-phenyl***	isomer 2: oil isomer 1: resin isomer 2: Harz
4.136	N	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	allyl***	
4.137	N	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	3-CF <sub>3</sub> -phenyl***	
4.138	N	NHCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	3-CF <sub>3</sub> -phenyl***	
4.139	N	OCH <sub>3</sub>	Me	-	1	2-CF <sub>3</sub>	oil
4.140	N	OCH <sub>3</sub>	Me	-	1	2-CH <sub>3</sub>	isomer 1: 105-107°C isomer 2: resin
4.141	N	NHCH <sub>3</sub>	Me	-	1	2-CF <sub>3</sub>	resin
4.142	N	NHCH <sub>3</sub>	Me	-	1	2-CH <sub>3</sub>	isomer 1: oil isomer 2: oil
4.143	CH	OCH <sub>3</sub>	CN	-	-	H	oil
4.144	CH	OCH <sub>3</sub>	CN	O	0	CH <sub>3</sub>	isomer 1: 136-138°C isomer 2: resin
4.145	CH	OCH <sub>3</sub>	CN	-	1	4-t-butyl	oil
4.146	CH	OCH <sub>3</sub>	CN	O	0	phenyl	oil
4.147	CH	OCH <sub>3</sub>	Me	O	0	CF <sub>2</sub> CHF <sub>2</sub>	isomer 1: 80-82°C
4.148	CH	OCH <sub>3</sub>	Me	O	0	CF <sub>2</sub> CHCl <sub>2</sub>	
4.149	CH	OCH <sub>3</sub>	Me	O	0	CF <sub>2</sub> CHBr <sub>2</sub>	
4.150	N	OCH <sub>3</sub>	Me	O	0	CF <sub>2</sub> CHF <sub>2</sub>	
4.151	N	OCH <sub>3</sub>	Me	O	0	CF <sub>2</sub> CHF <sub>2</sub>	
4.152	N	OCH <sub>3</sub>	Me	O	0	CF <sub>2</sub> CHF <sub>2</sub>	
4.153	N	NHCH <sub>3</sub>	Me	O	0	CF <sub>2</sub> CHF <sub>2</sub>	
4.154	N	NHCH <sub>3</sub>	Me	O	0	CF <sub>2</sub> CHF <sub>2</sub>	
4.155	CH	OCH <sub>3</sub>	Me	-	1	4-ethyl	isomer 1: oil isomer 2: oil
4.156	CH	OCH <sub>3</sub>	Me	-	1	4-t-butyl	isomer 1: resin isomer 2: oil
4.157	CH	OCH <sub>3</sub>	Me	-	1	4-CF <sub>3</sub>	isomer 1: resin isomer 2: 101-102°C
4.158	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	1	4-Cl-phenyl***	
4.159	CH	OMe	CN	O	0	phenyl	
4.160	N	OCH <sub>3</sub>	Me	-SO <sub>2</sub> -	0	Et	foam

- 50 -


Ex.	X	Y	R <sub>1</sub>	Z*	n	B or D	Physical data**
4.161	N	NHCH <sub>3</sub>	Me	-SO <sub>2</sub>	0	Et	141-143°C
4.162	CH	OCH <sub>3</sub>	Me	-OCH <sub>2</sub> -	0	SiMe <sub>3</sub>	isomer 1: resin isomer 2: oil
4.163	CH	OCH <sub>3</sub>	Me	O	0	CF <sub>2</sub> CHFCI	isomer 1: 75-76°C
4.164	CH	OCH <sub>3</sub>	Me	O	0	CF <sub>2</sub> CHFBr	isomer 1: resin
4.165	CH	OCH <sub>3</sub>	Me	O	0	CF <sub>2</sub> CHFCF <sub>3</sub>	isomer 1: 82-84°C
4.166	CH	OCH <sub>3</sub>	Me	S	0	Et	isomer 1: oil isomer 2: oil
4.167	CH	OCH <sub>3</sub>	Me	S	0	n-C <sub>3</sub> H <sub>7</sub>	isomer 1: oil isomer 2: oil
4.168	CH	OCH <sub>3</sub>	Me	-SCH <sub>2</sub> -	0	3-CF <sub>3</sub> -phenyl	
4.169	CH	OCH <sub>3</sub>	Me	-SO <sub>2</sub> CH <sub>2</sub> -	0	3-CF <sub>3</sub> -phenyl	
4.170	CH	OCH <sub>3</sub>	Me	-SO <sub>2</sub> CH <sub>2</sub> -	1	<u>B</u> 3-CF <sub>3</sub> -phenyl	<u>D</u> 3-F # #isom.1:112-114° #isomer 2: resin

\* The character "-" in column Z means the compound does not have a substituent -Z-B.

\*\* Physical data: m.p. or <sup>1</sup>H NMR of R<sub>1</sub> or R<sub>3</sub>, R<sub>3</sub> always being CH<sub>3</sub> in accordance with the structural formula in the heading of the table.

\*\*\* The radicals Z are to be read in such a manner that the skeleton of the structure is on the left of the structural element in question, while the substituent B is on the right. For example, the substituent Z-B in compound 4.107 is -CH<sub>2</sub>O-phenyl.

Tabelle 5



Ex. No.	X	Y	A	R <sub>1</sub>	R <sub>3</sub>	R <sub>6</sub>	R <sub>7</sub>	p	Physical data
5.1	CH	OCH <sub>3</sub>	NMe	Me	Me	Cl	Cl	1	
5.2	CH	OCH <sub>3</sub>	NMe	H	Me	Cl	Cl	1	
5.3	CH	OCH <sub>3</sub>	NMe	Me	Me	F	F	1	
5.4	CH	OCH <sub>3</sub>	NMe	Me	Et	F	F	1	
5.5	CH	OCH <sub>3</sub>	NMe	Me	Me	Br	Br	1	
5.6	CH	OCH <sub>3</sub>	NPh	Me	Me	Cl	Cl	1	
5.7	CH	OCH <sub>3</sub>	NPh	H	Me	Cl	Cl	1	
5.8	CH	OCH <sub>3</sub>	NPh	Me	Et	Cl	Cl	1	
5.9	CH	OCH <sub>3</sub>	O	Me	Me	Cl	Cl	1	isom.1: 86-88°C isom.2: resin isom.3: resin
5.10	CH	OCH <sub>3</sub>	O	H	Me	Cl	Cl	1	
5.11	CH	OCH <sub>3</sub>	O	Me	Me	F	F	1	
5.12	CH	OCH <sub>3</sub>	O	H	Me	F	F	1	
5.13	CH	OCH <sub>3</sub>	O	Me	H	F	F	1	
5.14	CH	OCH <sub>3</sub>	O	Me	C <sub>3</sub> H <sub>7</sub>	Cl	Cl	1	
5.15	CH	OCH <sub>3</sub>	O	Me	Δ	Cl	Cl	1	
5.16	CH	OCH <sub>3</sub>	O	Δ	Me	Cl	Cl	1	
5.17	CH	OCH <sub>3</sub>	O	Me	Δ	Cl	Cl	1	
5.18	CH	OCH <sub>3</sub>	O	Me	Me	Br	Br	1	foam
5.19	CH	OCH <sub>3</sub>	O	H	Me	Br	Br	1	
5.20	CH	OCH <sub>3</sub>	O	Me	Me	Cl	Cl	2	
5.21	CH	OCH <sub>3</sub>	O	H	Me	Cl	Cl	2	

- 52 -


Ex. No.	X	Y	A	R <sub>1</sub>	R <sub>3</sub>	R <sub>6</sub>	R <sub>7</sub>	p	Physical data
5.22	CH	OCH <sub>3</sub>	O	Me	Me	F	F	2	
5.23	N	OCH <sub>3</sub>	NMe	Me	Me	Cl	Cl	1	
5.24	N	OCH <sub>3</sub>	NMe	H	Me	Cl	Cl	1	
5.25	N	OCH <sub>3</sub>	NMe	Me	Me	F	F	1	
5.26	N	OCH <sub>3</sub>	NMe	Me	Et	F	F	1	
5.27	N	OCH <sub>3</sub>	NMe	Me		Br	Br	1	
5.28	N	OCH <sub>3</sub>	NPh	Me	Me	Cl	Cl	1	
5.29	N	OCH <sub>3</sub>	NPh	H	Me	Cl	Cl	1	
5.30	N	OCH <sub>3</sub>	NPh	Me	Et	Cl	Cl	1	
5.31	N	OCH <sub>3</sub>	O	Me	Me	Cl	Cl	1	
5.32	N	OCH <sub>3</sub>	O	H	Me	Cl	Cl	1	
5.33	N	OCH <sub>3</sub>	O	Me	Me	F	F	1	
5.34	N	OCH <sub>3</sub>	O	H	Me	F	F	1	
5.35	N	OCH <sub>3</sub>	O	Me	H	F	F	1	
5.36	N	OCH <sub>3</sub>	O	Me	C <sub>3</sub> H <sub>7</sub>	Cl	Cl	1	
5.37	N	OCH <sub>3</sub>	O	Me	Δ	Cl	Cl	1	
5.38	N	OCH <sub>3</sub>	O	Δ	Me	Cl	Cl	1	
5.39	N	OCH <sub>3</sub>	O	Me	Δ	Cl	Cl	1	
5.40	N	OCH <sub>3</sub>	O	Me	Me	Br	Br	1	
5.41	N	OCH <sub>3</sub>	O	H	Me	Br	Br	1	
5.42	N	OCH <sub>3</sub>	O	Me	Me	Cl	Cl	2	
5.43	N	OCH <sub>3</sub>	O	H	Me	Cl	Cl	2	
5.44	N	OCH <sub>3</sub>	O	Me	Me	F	F	2	
5.45	N	NHCH <sub>3</sub>	NMe	Me	Me	Cl	Cl	1	
5.46	N	NHCH <sub>3</sub>	NMe	H	Me	Cl	Cl	1	
5.47	N	NHCH <sub>3</sub>	NMe	Me	Me	F	F	1	
5.48	N	NHCH <sub>3</sub>	NMe	Me	Et	F	F	1	
5.49	N	NHCH <sub>3</sub>	NMe	Me		Br	Br	1	
5.50	CH	OCH <sub>3</sub>	O	Me	Me	H	H	1	isom.1: oil isom.2: oil
5.51	N	OCH <sub>3</sub>	O	Me	Me	Cl	Cl	1	isom.1: oil
5.52	N	NHCH <sub>3</sub>	O	Me	Me	Cl	Cl	1	137-139°C
5.53	N	NHCH <sub>3</sub>	O	Me	Me	Cl	Cl	1	

- 53 -


Ex. No.	X	Y	A	R <sub>1</sub>	R <sub>3</sub>	R <sub>6</sub>	R <sub>7</sub>	p	Physical data
---------	---	---	---	----------------	----------------	----------------	----------------	---	---------------

5.54	N	NHCH <sub>3</sub>	O	H	Me	Cl	Cl	1	
5.55	N	NHCH <sub>3</sub>	O	Me	Me	F	F	1	
5.56	N	NHCH <sub>3</sub>	O	H	Me	F	F	1	
5.57	N	NHCH <sub>3</sub>	O	Me	H	F	F	1	
5.58	N	NHCH <sub>3</sub>	O	Me	Me	Cl	Cl	2	
5.59	N	NHCH <sub>3</sub>	O	H	Me	Cl	Cl	2	
5.60	N	NHCH <sub>3</sub>	O	Me	Me	F	F	2	


and the compounds



isomer 1: 113-115°C / isomer 2: oil






isomer 1: oil / isomer 2: oil



resin

- 54 -




\*\* Physical data: m.p. or  $^1\text{H}$  NMR of  $\text{R}_1/\text{R}_2$  or  $\text{R}_3$




- 55 -

Table 6



Ex.	X	Y	R <sub>1</sub>	Z*	n	B or D	Physical data**
6.1	CH	OCH <sub>3</sub>	Me	O	0	allyl	oil
6.2	CH	OCH <sub>3</sub>	Me	-	1	4-Br	isom.1: oil isom.2: oil
6.3	N	OCH <sub>3</sub>	Me	O	0	allyl	oil
6.4	N	OCH <sub>3</sub>	Me	-	1	4-Br	131-132°C
6.5	N	NHCH <sub>3</sub>	Me	O	-	allyl	oil
6.6	N	NHCH <sub>3</sub>	Me	-	1	4-Br	isom.1: 117-119°C isom.2: oil
6.7	CH	OCH <sub>3</sub>	Me	-	1	4-methyl-	isomer 1: 2.10 isomer 2: 2.08
6.8	N	OCH <sub>3</sub>	Me	-	1	4-methyl-	
6.9	N	NHCH <sub>3</sub>	Me	-	1	4-methyl-	

Table 7:



Ex. No.	Y	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical d.**
7.1	OEt	NMe	Me	Me	Me	
7.2	OEt	NMe	H	Me	6-CF <sub>3</sub> -2-pyridyl	
7.3	OEt	NMe	Me	Δ	6-CF <sub>3</sub> -2-pyridyl	
7.4	OEt	NMe	Me	H	phenyl	
7.5	OEt	NMe	Me	Me	phenyl	
7.6	OEt	NMe	Δ	Me	phenyl	
7.7	OEt	NMe	H	Me	4-morpholinyl	
7.8	OEt	NPh	H	Me	phenyl	
7.9	OEt	NPh	Me	Me	phenyl	
7.10	OEt	NPh	Me	Δ	phenyl	
7.11	OEt	NPh	Δ	Me	phenyl	
7.12	OEt	O	Me	Me	Me	75°C
7.13	OEt	O	H	Me	Me	
7.14	OEt	O	Δ	Me	Me	
7.15	OEt	O	Me	Δ	Me	
7.16	OEt	O	Me	H	Me	
7.17	OEt	O	Me	Me	phenyl	
7.18	OEt	O	Me	Me	benzyl	
7.19	OEt	O	Me	Me	3-CF <sub>3</sub> -2-pyridyl	
7.20	OEt	O	Me	Me	3-CF <sub>2</sub> -phenyl	
7.21	NH <sub>2</sub>	NMe	Me	Δ	3-CF <sub>3</sub> -phenyl	
7.22	NH <sub>2</sub>	NMe	H	Me	4-CF <sub>3</sub> -phenyl	

- 57 -

Ex. No.	Y	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical d.**
7.23	NH <sub>2</sub>	NMe	Me	Me	4-CF <sub>3</sub> -phenyl	
7.24	NH <sub>2</sub>	NMe	Me	Me	2-chlorophenyl	
7.25	NH <sub>2</sub>	NMe	Me	Me	3-chlorophenyl	
7.26	NH <sub>2</sub>	NMe	Me	Me	4-chlorophenyl	
7.27	NH <sub>2</sub>	O	Me	Me	phenyl	
7.28	NH <sub>2</sub>	O	Me	Me	Me	
7.29	NH <sub>2</sub>	O	Me	Me	benzyl	
7.30	NH <sub>2</sub>	O	Me	Me	Et	
7.31	NH <sub>2</sub>	O	H	Me	Et	
7.32	NH <sub>2</sub>	O	Δ	Me	Et	
7.33	NH <sub>2</sub>	O	Me	Δ	Et	
7.34	NH <sub>2</sub>	O	Me	H	Et	
7.35	NH <sub>2</sub>	O	Me	Me	methoxymethyl	
7.36	NH <sub>2</sub>	O	Me	Δ	methoxymethyl	
7.37	NH <sub>2</sub>	O	Δ	Me	methoxymethyl	
7.38	NH <sub>2</sub>	O	Me	Me	ethoxymethyl	
7.39	NH <sub>2</sub>	O	H	Me	propargyl	
7.40	NH <sub>2</sub>	O	Me	Me	cyanomethyl	
7.41	NH <sub>2</sub>	O	Me	Me	H	
7.42	NH <sub>2</sub>	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub>	
7.43	NH <sub>2</sub>	O	Δ	Me	CF <sub>3</sub> CH <sub>2</sub>	
7.44	NH <sub>2</sub>	O	Me	H	CF <sub>3</sub> CH <sub>2</sub>	
7.45	NH <sub>2</sub>	O	Me	H	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	
7.46	NH <sub>2</sub>	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	
7.47	NH <sub>2</sub>	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	
7.48	NH <sub>2</sub>	O	Δ	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	
7.49	NHEt	NMe	Me	Δ	3-CF <sub>3</sub> -phenyl	
7.50	NHEt	NMe	H	Me	4-CF <sub>3</sub> -phenyl	
7.51	NHEt	NMe	Me	Me	4-CF <sub>3</sub> -phenyl	
7.52	NHEt	NMe	Me	Me	2-chlorophenyl	
7.53	NHEt	NMe	Me	Me	3-chlorophenyl	
7.54	NHEt	NMe	Me	Me	4-chlorophenyl	

- 58 -

Ex. No.	Y	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical d.**
7.55	NHEt	O	Me	Me	phenyl	77°C
7.56	NHEt	O	Me	Me	Me	
7.57	NHEt	O	Me	Me	benzyl	
7.58	NHEt	O	Me	Me	Et	
7.59	NHEt	O	H	Me	Et	
7.60	NHEt	O	Δ	Me	Et	
7.61	NHEt	O	Me	Δ	Et	
7.62	NHEt	O	Me	H	Et	
7.63	NHEt	O	Me	Me	methoxymethyl	
7.64	NHEt	O	Me	Δ	methoxymethyl	
7.65	NHEt	O	Δ	Me	methoxymethyl	
7.66	NHEt	O	Me	Me	ethoxymethyl	
7.67	NHEt	O	H	Me	propargyl	
7.68	NHEt	O	Me	Me	cyanomethyl	
7.69	NHEt	O	Me	Me	H	
7.70	NHEt	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub>	
7.71	NHEt	O	Δ	Me	CF <sub>3</sub> CH <sub>2</sub>	
7.72	NHEt	O	Me	H	CF <sub>3</sub> CH <sub>2</sub>	
7.73	NHEt	O	Me	H	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	
7.74	NHEt	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	
7.75	NHEt	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	
7.76	NHEt	O	Δ	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	
7.77	N(Me) <sub>2</sub>	NMe	Me	Δ	3-CF <sub>3</sub> -phenyl	
7.78	N(Me) <sub>2</sub>	NMe	H	Me	4-CF <sub>3</sub> -phenyl	
7.79	N(Me) <sub>2</sub>	NMe	Me	Me	4-CF <sub>3</sub> -phenyl	
7.80	N(Me) <sub>2</sub>	NMe	Me	Me	2-chlorophenyl	
7.81	N(Me) <sub>2</sub>	NMe	Me	Me	3-chlorophenyl	
7.82	N(Me) <sub>2</sub>	NMe	Me	Me	4-chlorophenyl	
7.83	N(Me) <sub>2</sub>	O	Me	Me	phenyl	
7.84	N(Me) <sub>2</sub>	O	Me	Me	Me	
7.85	N(Me) <sub>2</sub>	O	Me	Me	benzyl	
7.86	N(Me) <sub>2</sub>	O	Me	Me	Et	

- 59 -

Ex. No.	Y	A	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub> or NR <sub>3</sub> R <sub>4</sub>	Physical d.**
7.87	N(Me) <sub>2</sub>	O	H	Me	Et	
7.88	N(Me) <sub>2</sub>	O	Δ	Me	Et	
7.89	N(Me) <sub>2</sub>	O	Me	Δ	Et	
7.90	N(Me) <sub>2</sub>	O	Me	H	Et	
7.91	N(Me) <sub>2</sub>	O	Me	Me	methoxymethyl	
7.92	N(Me) <sub>2</sub>	O	Me	Δ	methoxymethyl	
7.93	N(Me) <sub>2</sub>	O	Δ	Me	methoxymethyl	
7.94	N(Me) <sub>2</sub>	O	Me	Me	ethoxymethyl	
7.95	N(Me) <sub>2</sub>	O	H	Me	propargyl	
7.96	N(Me) <sub>2</sub>	O	Me	Me	cyanomethyl	
7.97	N(Me) <sub>2</sub>	O	Me	Me	H	
7.98	N(Me) <sub>2</sub>	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub>	
7.99	N(Me) <sub>2</sub>	O	Δ	Me	CF <sub>3</sub> CH <sub>2</sub>	
7.100	N(Me) <sub>2</sub>	O	Me	H	CF <sub>3</sub> CH <sub>2</sub>	
7.101	N(Me) <sub>2</sub>	O	Me	H	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	
7.102	N(Me) <sub>2</sub>	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	
7.103	N(Me) <sub>2</sub>	O	Me	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	
7.104	N(Me) <sub>2</sub>	O	Δ	Me	CF <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub>	

## 2. Formulation Examples of active ingredients of the formula I (% = percent by weight)

### 2.1. Wettable powders

	a)	b)	c)
Active ingredient of Table 1-7	25 %	50 %	75 %
Sodium lignosulfonate	5 %	5 %	-
Sodium lauryl sulfate	3 %	-	5 %
Sodium diisobutyl naphthalenesulfonate	-	6 %	10 %
Octylphenol polyethylene glycol ether (7-8 mol of ethylene oxide)	-	2 %	-
Highly disperse silica	5 %	10 %	10 %
Kaolin	62 %	27 %	-

- 60 -

The active ingredient is mixed thoroughly with the additives, and the mixture is ground thoroughly in a suitable mill. This gives wettable powders which can be diluted with water to give suspensions of any desired concentration.

### 2.2. Emulsion concentrate

Active ingredient of Table 1-7	10 %
Octylphenol polyethylene glycol ether (4-5 mol of ethylene oxide)	3 %
Calcium dodecylbenzenesulfonate	3 %
Cyclohexanone	34 %
Xylene mixture	50 %

Emulsions of any desired dilution can be prepared from this concentrate by diluting it with water.

### 2.3. Dusts

	a)	b)
Active ingredient of Table 1-7	5 %	8 %
Talc	95 %	-
Kaolin	-	92 %

Ready-to-use dusts are obtained by mixing the active ingredient with the carrier and grinding the mixture in a suitable mill.

### 2.4. Extruder granules

Active ingredient of Table 1-7	10 %
Sodium lignosulfonate	2 %
Carboxymethylcellulose	1 %
Kaolin	87 %

The active ingredient is mixed with the additives, and the mixture is ground and moistened with water. This mixture is extruded and subsequently dried in a stream of air.

### 2.5. Coated granules

Active ingredient of Table 1-7	3 %
Polyethylene glycol (MW 200)	3 %
Kaolin	94 %

- 61 -

(MW = molecular weight)

In a mixer, the finely ground active ingredient is applied uniformly to the kaolin which has been moistened with polyethylene glycol. This gives dust-free coated granules.

#### 2.6. Suspension concentrate

Active ingredient of Table 1-7	40 %
Ethylene glycol	10 %
Nonylphenol polyethylene glycol ether (15 mol of ethylene oxid)	6 %
Sodium lignosulfonate	10 %
Carboxymethylcellulose	1 %
37% aqueous formaldehyde solution	0.2 %
Silicone oil in the form of a 75 % aqueous emulsion	0.8 %
Water	32 %

The finely ground active ingredient is mixed intimately with the additives. This gives a suspension concentrate from which suspensions of any desired dilution can be prepared by diluting it with water.

### 3. Biological Examples

#### A) Microbicidal action

##### Example B-1: Action against Phytophthora infestans on tomatoes

##### a) Curative action

Tomato plants cv. "Roter Gnom" are grown for three weeks and then sprayed with a zoospore suspension of the fungus and incubated in a cabinet at 18 to 20° and saturated atmospheric humidity. The humidification is interrupted after 24 hours. After the plants have dried, they are sprayed with a mixture which comprises the active ingredient, formulated as a wettable powder, at a concentration of 200 ppm. After the spray coating has dried, the plants are returned to the humid chamber for 4 days. Number and size of the typical leaf spots which have developed after this period are used to assess the activity of the test substances.

- 62 -

b) Preventive-systemic action

The active ingredient, formulated as a wettable powder, is applied at a concentration of 60 ppm (relative to the soil volume) to the soil surface of three-week old tomato plants cv. "Roter Gnom" in pots. After the plants have been left to stand for three days, the underside of the leaves is sprayed with a zoospore suspension of *Phytophthora infestans*. The plants are then kept for 5 days in a spray cabinet at 18 to 20°C and saturated atmospheric humidity. After this period, typical leaf spots develop, whose number and size are used for assessing the activity of the test substances.

While untreated, but infected, control plants show an infestation of 100 %, the active ingredients of the formula I in accordance with one of the tables, in particular the compounds No. 1.5, 1.24, 1.117, 2.24, 2.74 and 3.24, allow the infestation in both tests to be reduced to 20 % or less.

Example B-2: Action against *Plasmopara viticola* (Bert. et Curt.) (Berl. et De Toni) on vines

a) Residual-preventive action

Vine cuttings cv. "Chasselas" are grown in the greenhouse. When they have reached the 10-leaf stage, 3 plants are sprayed with a mixture (200 ppm of active ingredient). After the spray coating on the plants has dried, the leaf underside is inoculated uniformly with a spore suspension of the fungus. The plants are subsequently kept in a humid chamber for 8 days. After this period, the disease symptoms of the control plants are clearly developed. Number and size of the lesions on the treated plants are used to assess the activity of the test substances.

b) Curative action

Grape cuttings cv. "Chasselas" are grown in the greenhouse and, when they have reached the 10-leaf stage, are inoculated on the underside of the leaves with a spore suspension of *Plasmopara viticola*. After the plants have remained for 24 hours in a humid cabinet, they are sprayed with a mixture of active ingredient (200 ppm of active ingredient). The plants are subsequently kept for 7 more days in the humid cabinet. After this period, the disease symptoms have developed on the control plants. Number and size of the lesions on the treated plants are used to assess the activity of the test substances.

In comparison with the control plants, the infestation of plants which have been treated with active ingredients of the formula I is 20 % or less.



Example B-3: Action against *Pythium debaryanum* on sugar beet (*Beta vulgaris*)

a) Action after soil application

The fungus is grown on sterile oat grains and added to a mixture of soil and sand. This inoculated soil is filled into flower pots and sugar beet seeds are sown. Immediately after sowing, the test preparations, formulated as wettable powders, are poured over the soil in the form of an aqueous suspension (20 ppm of active ingredient relative to the soil volume). Thereupon, the pots are placed in a greenhouse for 2-3 weeks at 20-24°C. All the time, the soil is kept uniformly moist by spraying it gently with water. When the test is evaluated, the emergence of the sugar beet plants and the proportion of healthy and diseased plants is determined.

b) Action after seed-dressing

The fungus is grown on sterile oat grains and added to a mixture of soil and sand. This inoculated soil is filled into flower pots and sugar beet seeds which have been dressed with the test preparations, formulated as seed-dressing powders (1000 ppm of active ingredient relative to the weight of the seeds), are sown. The pots together with the seeds are placed in a greenhouse at 20-24°C for 2-3 weeks. The soil is kept uniformly moist by gently spraying it with water. When the test is evaluated, the emergence of the sugar beet plants and the proportion of healthy and diseased plants is determined.

After treatment with active ingredients of the formula I, in particular with the compounds No. 1.5, 1.24, 1.117, 2.24, 2.74 and 3.24, over 80 % of the plants emerge and have a healthy appearance. In the control pots, only few plants of unhealthy appearance have emerged.

Example B-4: Residual-protective action against *Cercospora arachidicola* on groundnuts

Groundnut plants 10 to 15 cm high are sprayed to drip point with an aqueous spray mixture (0.02 % of active ingredient) and, 48 hours later, inoculated with a conidia suspension of the fungus. The plants are incubated for 72 hours at 21° and high atmospheric humidity and subsequently placed in a greenhouse until the typical leaf spots have developed. The action of the active ingredient is assessed 12 days after the inoculation on the basis of the number and size of the leaf spots.

Active ingredients of the formula I cause a reduction in leaf spots to less than approximately 10 % of the leaf surface area. In some cases, the disease is reduced completely (0-5 % infestation).

Example B-5: Action against *Puccinia graminis* on wheat

- 64 -

a) Residual-protective action

6 days after sowing, wheat plants are sprayed to drip point with an aqueous spray mixture (0.02 % of active ingredient) and, 24 hours later, inoculated with a ureidospore suspension of the fungus. After an incubation time of 48 hours (conditions: relative humidity of 95 to 100 % at 20°), the plants are placed in a greenhouse at 22°. 12 days after the inoculation, the development of rust pustules is assessed.

b) Systemic action

5 days after sowing, an aqueous spray mixture (0.006 % of active ingredient relative to the soil volume) is poured next to wheat plants. Any contact of the spray mixture with aerial parts of the plants is carefully avoided. 48 hours later, the plants are inoculated with a ureidospore suspension of the fungus. After an incubation time of 48 hours (conditions: relative humidity of 95 to 100 % at 20°), the plants are placed in a greenhouse at 22°. 12 days after the inoculation, the development of rust pustules is assessed.

Compounds of the formula I, in particular those of Table 1, in particular No. 1.5, 1.24, 1.117, 2.74, cause a marked reduction in fungus infestation, in some cases down to 10-0 %.

Example B-6: Action against *Pyricularia oryzae* on ricea) Residual-protective action

Rice plants are grown for two weeks and then sprayed to drip point with an aqueous spray mixture (0.02 % of active ingredient), and, 48 hours later, inoculated with a conidia suspension of the fungus. The fungus infestation is assessed 5 days after the inoculation, during which period the relative atmospheric humidity was kept at 95 to 100 % and the temperature at 22°.

b) Systemic action

An aqueous spray mixture (0.006 % of active ingredient relative to the soil volume) is poured next to 2-week old rice plants. Contact of the spray mixture with the aerial parts of the plants is carefully avoided. The pots are then filled with such an amount of water that the lowest parts of the stems of the rice plants are submerged. After 96 hours, the plants are inoculated with a conidia suspension of the fungus and kept for 5 days at a relative atmospheric humidity of 95 to 100 % and a temperature of 24°C.

In many cases, compounds of the formula I prevent the development of disease on the infected plants.

- 65 -

Example B-7: Residual-protective action against *Venturia inaequalis* on apples

Apple cuttings having fresh shoots 10 to 20 cm in length are sprayed to drip point with a spray mixture (0.02 % of active ingredient) and, 24 hours later, inoculated with a conidia suspension of the fungus. The plants are incubated for 5 days at a relative atmospheric humidity of 90 to 100 % and kept in a greenhouse for a further 10 days at 20 to 24°. The infestation with scab was assessed 15 days after the inoculation.

Most of the compounds of the formula I of one of Tables 1, 2 or 3 have a sustainable action against scab diseases.

Example B-8: Action against *Erysiphe graminis* on barley

a) Residual-protective action

Barley plants approximately 8 cm high are sprayed to drip point with an aqueous spray mixture (0.02 % of active ingredient) and, 3 to 4 hours later, dusted with conidia of the fungus. The inoculated plants are placed in a greenhouse at 22°. The fungus infestation is assessed 10 days after the inoculation.

b) Systemic action

An aqueous spray mixture (0.002 % of active ingredient relative to the soil volume) is poured next to barley plants approximately 8 cm high. Contact of the spray mixture with aerial parts of the plants is carefully avoided. 48 hours later, the plants are dusted with conidia of the fungus. The inoculated plants are placed in a greenhouse at 22°. The fungus infestation is assessed 10 days after the inoculation.

Compounds of the formula I, in particular compounds No. 1.117, 2.24 and 3.24, are generally capable of reducing the disease to less than 20 %, in some cases also completely.

Example B-9: Action against *Podosphaera leucotricha* on apple shoots

Residual-protective action

Apple cuttings having fresh shoots approximately 15 cm long are sprayed with a spray mixture (0.06 % of active ingredient). After 24 hours, the treated plants are inoculated with a conidia suspension of the fungus and placed in a controlled-environment cabinet at a relative atmospheric humidity of 70 % and at 20°C. The fungus infestation is assessed 12 days after the inoculation.

Active ingredients of the formula I allow the disease to be reduced to less than 20 %. Control plants show a disease level of 100 %.

Example B-10: Action against *Botrytis cinerea* on apple fruits. Residual- protective action

- 66 -

Artificially wounded apples are treated by applying dropwise a spray mixture (0.02 % of active ingredient) to the wounds. The treated fruits are subsequently inoculated with a spore suspension of the fungus and incubated for one week at high atmospheric humidity and at approximately 20°C. The fungicidal action of the test substance is deduced from the number of wounds showing signs of rot.

Active ingredients of the formula I are capable of reducing the spread of rot, in some cases completely.

Example B-11: Action against *Helminthosporium gramineum*

Wheat grains are contaminated with a spore suspension of the fungus and allowed to dry. The contaminated grains are dressed with a suspension of the test substance (600 ppm of active ingredient relative to the weight of the seeds). After two days, the grains are arranged in suitable dishes containing agar, and, after a further four days, the development of fungal colonies around the grains is assessed. The test substance is assessed on the basis of the number and size of the fungal colonies.

A good action, i.e. inhibition of fungal colonies, is shown, in some cases, by compounds of the formula I.

Example B-12: Action against *Colletotrichum lagenarium* on cucumbers

Cucumber plants are grown for two weeks and then sprayed with a spray mixture (concentration 0.002 %). After 2 days, the plants are inoculated with a spore suspension ( $1.5 \times 10^5$  spores/ml) of the fungus and incubated for 36 hours at 23° and high atmospheric humidity. The incubation is then continued at normal atmospheric humidity at approximately 22-23°C. The fungus infestation which has developed is assessed 8 days after the inoculation. Untreated, but infected, control plants show a fungus infestation of 100 %.

A virtually complete inhibition of the infestation with disease is shown, in some cases, by compounds of the formula I, in particular by No. 1.5 and 3.24.

Example B-13: Action against *Fusarium nivale* on rye

Rye cv. Tetrahell, naturally infected with *Fusarium nivale*, is dressed with the test fungicide using a mixing roller, the following concentrations being used: 20 or 6 ppm of a.i. (relative to the weight of the seed).

Using a seed drill, the infected and treated rye is drilled in October in a field in 3 m plots of 6 seed furrows. 3 replicates per concentration.

- 67 -

Until the infestation is evaluated, the test plants are grown under normal field conditions (preferably in a region with a complete covering of snow during the winter months).

To assess the phytotoxicity, seed emergence is scored in autumn and number of plants per unit area/number of tillers per plant in spring.

To determine the activity of the active ingredient, the percentage of Fusarium-infested plants is counted immediately after the snow has melted. In the present case, the number of infested plants was lower than 5 %. The plants which emerged had a healthy appearance.

Example B-14: Action against Septoria nodorum on wheat

Wheat plants in the 3-leaf stage are sprayed with a spray mixture (60 ppm of a.i.) prepared with a wettable powder of the active ingredients (2.8:1).

After 24 hours the treated plants are inoculated with a conidia suspension of the fungus. The plants are subsequently incubated for 2 days at a relative atmospheric humidity of 90-100 % and placed in a greenhouse at 20-24°C for a further 10 days. 13 days after the inoculation, the fungus infestation is assessed. Less than 1 % of the wheat plants were infested.

Example B-15: Action against Rhizoctonia solani on rice

Protective-local soil application

The soil around 10-day old rice plants was watered with a suspension (spray mixture) prepared with a formulated test substance without aerial parts of the plants coming into contact with the suspension. Three days later, the plants were inoculated by placing a stem of barley straw infected with *Rhizoctonia solani* between the rice plants of each pot. After incubation for 6 days in a controlled-environment cabinet at 29°C day-time and 26°C night-time temperature and a relative atmospheric humidity of 95 %, the fungus infestation is assessed. Fewer than 5 % of the rice plants are infested. The plants had a healthy appearance.

Protective-local foliar application

12-day old rice plants are sprayed with a suspension prepared with formulated test substances. One day later, inoculation is carried out by placing a stem of barley straw,

- 68 -

infected with *Rhizoctonia solani*, between the rice plants of each pot. The plants were scored after incubation for 6 days in a controlled-environment cabinet at 29°C day-time and 26°C night-time temperature and a relative atmospheric humidity of 95 %. Untreated, but inoculated, control plants showed a fungal infestation of 100 %. Compounds of the formula I cause in some cases complete inhibition of the infestation with disease.

#### B. Insecticidal action

##### Example B-16: Action against *Aphis craccivora*

Pea seedlings are populated with *Aphis craccivora*, subsequently sprayed with a spray mixture comprising 400 ppm of active ingredient and then incubated at 20°. The percentage reduction in population (% action) is determined after 3 and 6 days by comparing the number of dead aphids on the treated and the untreated plants. In this test, compounds of Tables 1-7 have a good action. Compound No. 4.5, isomer 1, in particular, has an action of over 80 %.

##### Example B-17: Action against *Diabrotica balteata*

Maize seedlings are sprayed with an aqueous emulsion spray mixture comprising 400 ppm of active ingredient, then, after the spray coating has dried on, populated with 10 second instar larvae of *Diabrotica balteata* and subsequently placed in a plastic container. The percentage reduction in population (% action) is determined after 6 days by comparing the number of dead larvae of the treated and untreated plants.

In this test, compounds of Tables 1-7 have a good action. In particular, compounds No. 1.5, 1.24, 2.24, 3.24, 4.5, isomer 1, 4.14, isomer 1, 4.14, isomer 1, 4.117, both isomers, 4.121, 4.170, 5.9, isomer 1 and 5.51, isomer 1, have an action of over 80 %.

##### Example B-18: Action against *Heliothis virescens*

Young soya bean plants are sprayed with an aqueous emulsion spray mixture comprising 400 ppm of active ingredient, then, after the spray coating has dried on, populated with 10 first instar caterpillars of *Heliothis virescens* and subsequently placed in a plastic container. The percentage reduction in population and the feeding damage (% action) are determined after 6 days by comparing the number of dead caterpillars and the feeding damage of the treated and untreated plants.

In this test, compounds of Tables 1-7 show a good action. In particular, compounds No. 4.5, isomer 1, 4.14, isomer 1, 4.18, 4.117, both isomers, 4.121, 5.9, isomer 1, and 5.52, isomer 1, have an action of over 80 %.

Example B-19: Action against *Spodoptera littoralis*

Young soya bean plants are sprayed with an aqueous emulsion spray mixture comprising 400 ppm of active ingredient, then, after the spray coating has dried on, populated with 10 third instar caterpillars of *Spodoptera littoralis* and subsequently placed in a plastic container. The percentage reduction in population and the percentage reduction in feeding damage (% action) are determined after 3 days by comparing the number of dead caterpillars and the feeding damage of the treated and untreated plants.

In this test, compounds of Tables 1-7 show a good action. In particular, compounds No. 1.5, 1.24, 2.24, 3.24, 4.18, 4.5, isomer 1, 5.9, isomer 1, and 5.52, isomer 1, have an action of over 80 %.

C. Acaricidal activityExample B-20: Action against *Tetranychus urticae*

Young bean plants are populated with a mixed population of *Tetranychus urticae*, then, 1 day later, sprayed with an aqueous emulsion spray mixture comprising 400 ppm of active ingredient, incubated for 6 days at 25° and subsequently evaluated. The percentage reduction in population (% action) is determined by comparing the number of dead eggs, larvae and adults on the treated and untreated plants.

In this test, compounds of Tables 1-6 show a good action. In particular, compounds No. 1.5, 1.24, 2.24 und 3.24, 4.5, isomer 1, 4.14, isomer 1, 4.117, both isomers, 5.9, isomer 1, and 5.52, isomer 1, have an action of over 80 %.


Example B-21: Action against *Boophilus microplus*

Adult female ticks which have sucked themselves full are attached to a PVC board and covered with a cottonwool ball, and 10 ml of aqueous test solution comprising 125 ppm of active ingredient are poured over the ticks. The cottonwool ball is removed, and the ticks are incubated for 4 weeks for oviposition. The action becomes apparent either, in the case of the female, in the form of mortality or sterility or, in the case of the eggs, in the form of an ovicidal action.

- 70 -

What is claimed is:

1. A compound of the formula I



and the isomers, and isomer mixtures, thereof which are possible, in which either

- a) X is an N atom and  
Y is OR<sub>11</sub> or N(R<sub>12</sub>)R<sub>13</sub>, or
- b) X is CH and  
Y is OR<sub>11</sub>,


and in which furthermore:


R<sub>11</sub> is C<sub>1</sub>-C<sub>4</sub>alkyl;

R<sub>12</sub> and R<sub>13</sub>, independently, are hydrogen or C<sub>1</sub>-C<sub>4</sub>alkyl;

A is an O atom or the group NR<sub>4</sub>;

R<sub>1</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, halo-C<sub>1</sub>-C<sub>4</sub>alkyl, cyclopropyl, cyano or methylthio;

R<sub>2</sub> is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, a group , or

a group , or thienyl;

D radicals are identical or different and are halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy,

C<sub>1</sub>-C<sub>2</sub>haloalkyl, C<sub>1</sub>-C<sub>2</sub>haloalkoxy, C<sub>3</sub>-C<sub>6</sub>alkenyloxy, C<sub>3</sub>-C<sub>6</sub>alkynyloxy,

C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, cyano or nitro;

n is 0, 1, 2, 3 or 4;

Z is -O-, -O-(C<sub>1</sub>-C<sub>4</sub>alkyl)-, -(C<sub>1</sub>-C<sub>4</sub>alkyl)-O-, -S(O)<sub>m</sub>-, -(C<sub>1</sub>-C<sub>4</sub>alkyl)-S(O)<sub>m</sub>-,

-S(O)<sub>m</sub>-(C<sub>1</sub>-C<sub>4</sub>alkyl)-,

m is 0, 1 or 2,


B is C<sub>1</sub>-C<sub>6</sub>alkyl, halo-C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, or is C<sub>2</sub>-C<sub>6</sub>alkenyl or

C<sub>2</sub>-C<sub>4</sub>alkynyl-C<sub>1</sub>-C<sub>2</sub>alkyl, each of which is unsubstituted or substituted by 1 to 3 halogen



- 71 -

atoms, or is aryl or heterocyclyl, each of these two, independently, being unsubstituted or mono- to pentasubstituted by C<sub>1</sub>-C<sub>6</sub>alkyl, halo-C<sub>1</sub>-C<sub>6</sub>alkyl, halogen, C<sub>1</sub>-C<sub>6</sub>alkoxy or

halo-C<sub>1</sub>-C<sub>6</sub>alkoxy, or a group , or trimethylsilyl;

R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub> and R<sub>9</sub>, independently, hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl or halogen and p is 0, 1, 2 or 3;

R<sub>3</sub> is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, C<sub>1</sub>-C<sub>6</sub>haloalkyl having 1 to 5 halogen atoms, C<sub>1</sub>-C<sub>4</sub>alkoxy-C<sub>1</sub>-C<sub>2</sub>alkyl, C<sub>2</sub>-C<sub>4</sub>alkenyl-C<sub>1</sub>-C<sub>2</sub>alkyl, which is unsubstituted or substituted by 1 to 3 halogen atoms, C<sub>2</sub>-C<sub>4</sub>alkynyl-C<sub>1</sub>-C<sub>2</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl which is unsubstituted or substituted by 1 to 4 halogen atoms, C<sub>3</sub>-C<sub>6</sub>-cycloalkyl-C<sub>1</sub>-C<sub>4</sub>alkyl which is unsubstituted or substituted by 1 to 4 halogen atoms, cyano-C<sub>1</sub>-C<sub>4</sub>alkyl; C<sub>1</sub>-C<sub>4</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>2</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>2</sub>alkyl, phenyl-C<sub>1</sub>-C<sub>3</sub>alkyl which is unsubstituted or substituted by halogen, C<sub>1</sub>-C<sub>3</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>haloalkyl, cyano, nitro or C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, it being possible for the phenyl group to be monosubstituted to trisubstituted by identical or different substituents; phenyl which is unsubstituted or mono- to disubstituted, independently, by C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, halogen, C<sub>1</sub>-C<sub>2</sub>haloalkyl having 1 to 3 halogen atoms, nitro or cyano, or pyridyl which is unsubstituted or mono- to disubstituted, independently, by C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, halogen, C<sub>1</sub>-C<sub>2</sub>haloalkyl having 1 to 3 halogen atoms, nitro or cyano;

R<sub>4</sub> is C<sub>1</sub>-C<sub>4</sub>alkyl, phenyl, or

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 7-membered ring which is unsubstituted or substituted by C<sub>1</sub>-C<sub>4</sub>alkyl and which can have 1 to 3 additional hetero atoms selected from amongst N, O and S.

2. A compound of the formula I according to claim 1, in which either


- a) X is an N atom and  
Y is OCH<sub>3</sub> or NHCH<sub>3</sub>, or
- b) X is CH and  
Y is OCH<sub>3</sub>,


and in which furthermore,

A is an O atom or the group NR<sub>4</sub>;

R<sub>1</sub> is hydrogen, C<sub>1</sub>-C<sub>4</sub>alkyl, halo-C<sub>1</sub>-C<sub>4</sub>alkyl, cyclopropyl, cyano or methylthio;

- 72 -

$R_2$  is hydrogen,  $C_1$ - $C_6$ alkyl,  $C_3$ - $C_6$ cycloalkyl, a group , a group



D radicals are identical or different and are halogen,  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ alkoxy,  $C_1$ - $C_2$ haloalkyl,  $C_1$ - $C_2$ haloalkoxy,  $C_3$ - $C_6$ alkenyloxy,  $C_3$ - $C_6$ alkynyloxy,  $C_1$ - $C_4$ alkylenedioxy, cyano or nitro;


n is 0, 1, 2, 3 or 4;

Z is -O-, -O-( $C_1$ - $C_4$ alkyl)-, -( $C_1$ - $C_4$ alkyl)-O-, -S(O)<sub>m</sub>-, -( $C_1$ - $C_4$ alkyl)-S(O)<sub>m</sub>-, -S(O)<sub>m</sub>-( $C_1$ - $C_4$ alkyl)-,

m is 0, 1 or 2,

B is  $C_1$ - $C_6$ alkyl, halo- $C_1$ - $C_6$ alkyl,  $C_3$ - $C_6$ cycloalkyl,

or is  $C_2$ - $C_6$ alkenyl or  $C_2$ - $C_4$ alkynyl- $C_1$ - $C_2$ alkyl, each of which is unsubstituted or substituted by 1 to 3 halogen atoms, or is aryl or heterocyclyl, or aryl or heterocyclyl, each of these, independently, being monosubstituted to pentasubstituted by  $C_1$ - $C_6$ alkyl, halo- $C_1$ - $C_6$ alkyl, halogen,  $C_1$ - $C_6$ alkoxy or halo- $C_1$ - $C_6$ alkoxy, or a group



$R_5$ ,  $R_6$ ,  $R_7$ ,  $R_8$  and  $R_9$ , independently, are hydrogen,  $C_1$ - $C_4$ alkyl or halogen and p is 0, 1, 2 or 3;

$R_3$  is hydrogen,  $C_1$ - $C_6$ alkyl,  $C_1$ - $C_6$ haloalkyl having 1 to 5 halogen atoms,  $C_1$ - $C_4$ alkoxy- $C_1$ - $C_2$ alkyl,  $C_2$ - $C_4$ alkenyl- $C_1$ - $C_2$ alkyl which is unsubstituted or substituted by 1 to 3 halogen atoms,  $C_2$ - $C_4$ alkynyl- $C_1$ - $C_2$ alkyl,  $C_3$ - $C_6$ cycloalkyl which is unsubstituted or substituted by 1 to 4 halogen atoms,  $C_3$ - $C_6$ cycloalkyl- $C_1$ - $C_4$ alkyl which is unsubstituted or substituted by 1 to 4 halogen atoms, cyano- $C_1$ - $C_4$ alkyl;  $C_1$ - $C_4$ alkoxycarbonyl- $C_1$ - $C_2$ alkyl, phenyl- $C_1$ - $C_3$ alkyl which is unsubstituted or substituted by halogen,  $C_1$ - $C_3$ alkyl,  $C_1$ - $C_4$ alkoxy,  $C_1$ - $C_4$ haloalkyl, cyano, nitro,  $C_1$ - $C_4$ alkylenedioxy, it being possible for the phenyl group to be monosubstituted to trisubstituted by identical or different substituents; phenyl which is unsubstituted or mono- to disubstituted, independently, by  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ alkoxy, halogen,  $C_1$ - $C_2$ haloalkyl having 1 to 3 halogen atoms, nitro or cyano, or pyridyl which is unsubstituted or mono- to disubstituted, independently, by  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ alkoxy, halogen,  $C_1$ - $C_2$ haloalkyl having 1 to 3

- 73 -

halogen atoms, nitro or cyano;

R<sub>4</sub> is C<sub>1</sub>-C<sub>4</sub>alkyl or phenyl, or

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 7-membered ring which is unsubstituted or substituted by C<sub>1</sub>-C<sub>4</sub>alkyl and which can have 1 to 3 additional hetero atoms selected from amongst N, O and S.

3. A compound of the formula I according to claim 2, in which

X is N and

Y is OCH<sub>3</sub>.

4. A compound of the formula I according to claim 2, in which

X is CH.

5. A compound of the formula I according to claim 2, in which

X is N,

Y is NHCH<sub>3</sub> and

R<sub>1</sub> is H, CH<sub>3</sub>, cyclopropyl or CN.

6. A compound of the formula I according to any one of claims 2 to 5, in which

A is oxygen, NCH<sub>3</sub> or n-C<sub>6</sub>H<sub>5</sub>.


7. A compound of the formula I according to any one of claims 2 to 6, in which

R<sub>1</sub> is hydrogen, methyl, cyclopropyl or cyano.

8. A compound of the formula I according to any one of claims 2 to 7, in which

R<sub>2</sub> is C<sub>1</sub>-C<sub>4</sub>alkyl or cyclopropyl.


9. A compound of the formula I according to any one of claims 2 to 6, in which

R<sub>2</sub> is a group  and

D is halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>2</sub>alkyl which is substituted by 1 to 5 halogen atoms, C<sub>1</sub>-C<sub>2</sub>haloalkoxy, C<sub>3</sub>-C<sub>6</sub>alkenyloxy, C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, cyano, or nitro, or thienyl, and  
n is 0, 1, 2, 3 or 4.


- 74 -

10. A compound of the formula I according to any one of claims 2 to 7, in which


$R_2$  is a group  and

Z is -O-, -O-(C<sub>1</sub>-C<sub>4</sub>alkyl)-, -(C<sub>1</sub>-C<sub>4</sub>alkyl)-O-, -S(O)<sub>2</sub>-, -(C<sub>1</sub>-C<sub>4</sub>alkyl)-S(O)<sub>2</sub>-,  
-S(O)<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub>alkyl)-.

11. A compound of the formula I according to any one of claims 2 to 7, in which

$R_2$  is a group  and

B is C<sub>1</sub>-C<sub>4</sub>alkyl, halogen-C<sub>1</sub>-C<sub>4</sub>alkyl, or is C<sub>2</sub>-C<sub>4</sub>alkenyl or C<sub>2</sub>-C<sub>4</sub>alkynyl-C<sub>1</sub>-C<sub>2</sub>alkyl, each of which is unsubstituted or substituted by 1 to 3 halogen atoms, or is aryl or aryl which is monosubstituted or disubstituted, independently, by C<sub>1</sub>-C<sub>2</sub>alkyl, halo-C<sub>1</sub>-C<sub>2</sub>alkyl, halogen,

C<sub>1</sub>-C<sub>2</sub>alkoxy or halo-C<sub>1</sub>-C<sub>2</sub>alkoxy, or is a group 

$R_5$ ,  $R_6$ ,  $R_7$ ,  $R_8$  and  $R_9$ , independently, are hydrogen, C<sub>1</sub>-C<sub>2</sub>alkyl or halogen, and p is 0, 1, 2 or 3.

12. A compound of the formula I according to any one of claims 2 to 7, in which  $R_2$  is a phenyl group which is substituted in the 4-position by -Z-B.

13. A compound of the formula I according to any one of claims 2 to 12, in which  $R_3$  is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl or C<sub>1</sub>-C<sub>4</sub>haloalkyl having 1 to 3 halogen atoms, C<sub>1</sub>-C<sub>2</sub>alkoxy-C<sub>1</sub>-C<sub>2</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>2</sub>alkyl, propenyl which is unsubstituted or substituted by 1 to 3 halogen atoms, propargyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, cyclopropylmethyl which is unsubstituted or substituted by 1 to 2 halogen atoms, cyano-C<sub>1</sub>-C<sub>2</sub>alkyl, phenyl-C<sub>1</sub>-C<sub>2</sub>alkyl which is unsubstituted or substituted by halogen, methyl, methoxy or halomethyl having 1 to 3 halogen atoms, it being possible for the phenyl group to be monosubstituted to disubstituted by identical or different substituents; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or pyridyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or

- 75 -

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 7-membered ring which is unsubstituted or substituted by C<sub>1</sub>-C<sub>4</sub>alkyl and which can have 1 to 3 additional hetero atoms selected from amongst N, O and S.

14. A compound of the formula I according to any one of claims 2 to 13, in which R<sub>4</sub> is methyl or phenyl.

15. A compound of the formula I according to any one of claims 1 to 4, in which

X is CH;

Y is OCH<sub>3</sub>;

R<sub>1</sub> is CH<sub>3</sub>;

A is oxygen;

R<sub>2</sub> is 4-methylphenyl or 4-allyloxyphenyl or 4-(3-trifluoromethyl-benzyloxy)phenyl or 4-(2,2-dichlorocyclopropylmethoxy)phenyl and

R<sub>3</sub> is CH<sub>3</sub>.

16. A compound of the formula I according to claim 1, in which either

a) X is an N atom and

Y is OCH<sub>3</sub> or NHCH<sub>3</sub>, or

b) X is CH and

Y is OCH<sub>3</sub>,

and in which furthermore

A is an O atom or the group NR<sub>4</sub>;

R<sub>1</sub> is hydrogen; C<sub>1</sub>-C<sub>4</sub>alkyl; cyclopropyl; cyano or methylthio;

R<sub>2</sub> is hydrogen; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>3</sub>-C<sub>6</sub>cycloalkyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>2</sub>haloalkyl, C<sub>1</sub>-C<sub>2</sub>haloalkoxy, C<sub>3</sub>-C<sub>6</sub>alkenyloxy, C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, cyano or nitro; thienyl;

R<sub>3</sub> is hydrogen; C<sub>1</sub>-C<sub>6</sub>alkyl; C<sub>1</sub>-C<sub>6</sub>haloalkyl having 1 to 5 halogen atoms;

C<sub>1</sub>-C<sub>4</sub>alkoxy-C<sub>1</sub>-C<sub>2</sub>alkyl; C<sub>2</sub>-C<sub>4</sub>alkenyl-C<sub>1</sub>-C<sub>2</sub>alkyl which is unsubstituted or substituted by 1 to 3 halogen atoms; C<sub>2</sub>-C<sub>4</sub>alkynyl-C<sub>1</sub>-C<sub>2</sub>alkyl; C<sub>3</sub>-C<sub>6</sub>cycloalkyl which is unsubstituted or substituted by 1 to 4 halogen atoms; C<sub>3</sub>-C<sub>6</sub>cycloalkyl-C<sub>1</sub>-C<sub>4</sub>alkyl which is unsubstituted or substituted by 1 to 4 halogen atoms; cyano-C<sub>1</sub>-C<sub>4</sub>alkyl;

C<sub>1</sub>-C<sub>4</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>2</sub>alkyl; phenyl-C<sub>1</sub>-C<sub>3</sub>alkyl which is unsubstituted or substituted by halogen, C<sub>1</sub>-C<sub>3</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>haloalkyl, cyano, nitro, C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, it being possible for the phenyl group to be monosubstituted to trisubstituted by identical

- 76 -

or different substituents; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, halogen, C<sub>1</sub>-C<sub>2</sub>haloalkyl having 1 to 3 halogen atoms, nitro or cyano; pyridyl which is unsubstituted or monosubstituted to disubstituted, independently, by C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, halogen, C<sub>1</sub>-C<sub>2</sub>haloalkyl having 1 to 3 halogen atoms, nitro or cyano;

R<sub>4</sub> is C<sub>1</sub>-C<sub>4</sub>alkyl; phenyl; or

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 7-membered ring which is unsubstituted or substituted by C<sub>1</sub>-C<sub>4</sub>alkyl and which can have 1 to 3 additional hetero atoms selected from amongst N, O and S.

17. A compound of the formula I according to claim 16, in which

X is CH or N

Y is OCH<sub>3</sub>

A is O or N-R<sub>4</sub>

R<sub>1</sub> is methyl, cyclopropyl or methylthio;

R<sub>2</sub> is methyl; cyclopropyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>2</sub>alkyl which is substituted by 1 to 5 halogen atoms, C<sub>1</sub>-C<sub>2</sub>haloalkoxy, C<sub>3</sub>-C<sub>6</sub>-alkenyloxy,

C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, cyano, or nitro; or thienyl; and in which

R<sub>3</sub> is as described in formula I and

R<sub>4</sub> is methyl or phenyl, or

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded are either pyrrolidine, piperidine, morpholine, thiomorpholine, hexamethyleneimine, imidazole, pyrazole, pyrrole, 1,2,4-triazole or 1,2,3-triazole.

18. A compound of the formula I according to claim 16, in which

X is N,

Y is NHCH<sub>3</sub>,

A is O or N-R<sub>4</sub>,

R<sub>1</sub> is methyl, cyclopropyl or methylthio;

R<sub>2</sub> is methyl; cyclopropyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>2</sub>alkyl which is substituted by 1 to 5 halogen atoms, C<sub>1</sub>-C<sub>2</sub>haloalkoxy, C<sub>3</sub>-C<sub>6</sub>alkenyloxy,

C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, cyano or nitro; or thienyl; and in which

R<sub>3</sub> is as described in formula I and

R<sub>4</sub> is methyl or phenyl, or

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded are either pyrrolidine, piperidine, morpholine, thiomorpholine, hexamethyleneimine, imidazole, pyrazole, pyrrole, 1,2,4-triazole or 1,2,3-triazole.

19. A compound of the formula I according to claim 16, in which:

A is an O atom,

R<sub>1</sub> is methyl,

R<sub>2</sub> is methyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>2</sub>alkyl which is substituted by 1 to 5 halogen atoms, C<sub>1</sub>-C<sub>2</sub>haloalkoxy, C<sub>3</sub>-C<sub>6</sub>alkenyloxy, C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, cyano or nitro; or thienyl; and

R<sub>3</sub> is C<sub>1</sub>-C<sub>6</sub>alkyl.

20. A compound of the formula I according to claim 16, in which:

R<sub>1</sub> is methyl,

R<sub>2</sub> is methyl and

R<sub>3</sub> is as described in claim 1 and

R<sub>4</sub> is methyl or phenyl, or

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 7-membered ring which is unsubstituted or substituted by C<sub>1</sub>-C<sub>4</sub>alkyl and which can have 1 to 3 additional hetero atoms selected from amongst N, O and S.

21. A compound of the formula I according to claim 16, in which:

R<sub>3</sub> is hydrogen; C<sub>1</sub>-C<sub>4</sub>alkyl; C<sub>1</sub>-C<sub>4</sub>haloalkyl having 1 to 3 halogen atoms;

C<sub>1</sub>-C<sub>2</sub>alkoxy-C<sub>1</sub>-C<sub>2</sub>alkyl; propenyl which is unsubstituted or substituted by 1 to 3 halogen atoms; propargyl; C<sub>3</sub>-C<sub>6</sub>cycloalkyl; C<sub>3</sub>-C<sub>6</sub>cycloalkylmethyl which is unsubstituted or substituted by 1 to 2 halogen atoms; cyano-C<sub>1</sub>-C<sub>2</sub>alkyl;

C<sub>1</sub>-C<sub>2</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>2</sub>alkyl; phenyl-C<sub>1</sub>-C<sub>2</sub>alkyl which is unsubstituted or substituted by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano, nitro or C<sub>1</sub>-C<sub>2</sub>alkylenedioxy, it being possible for the phenyl group to be monosubstituted to disubstituted by identical or different substituents; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or pyridyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro;

R<sub>4</sub> is methyl or phenyl; or

- 78 -

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded form a saturated or unsaturated 5- to 7-membered ring which is unsubstituted or substituted by C<sub>1</sub>-C<sub>4</sub>alkyl and which can have 1 to 3 additional hetero atoms selected from amongst N, O and S.

22. A compound of the formula I according to claim 21, in which

R<sub>3</sub> and R<sub>4</sub> together with the nitrogen atom to which they are bonded are triazolyl, morpholinyl, 2,6-dimethylmorpholinyl, azepinyl, piperidyl or pyrrolidinyl.

23. A compound of the formula I according to claim 16, in which:

A is an O atom,

R<sub>1</sub> is methyl,

R<sub>2</sub> is methyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>2</sub>alkyl which is substituted by 1 to 5 halogen atoms, C<sub>1</sub>-C<sub>2</sub>haloalkoxy, C<sub>3</sub>-C<sub>6</sub>alkenyloxy, C<sub>3</sub>-C<sub>6</sub>alkynyloxy, C<sub>1</sub>-C<sub>4</sub>alkylenedioxy, cyano or nitro; or thienyl; and

R<sub>3</sub> is hydrogen; C<sub>1</sub>-C<sub>4</sub>alkyl; C<sub>1</sub>-C<sub>4</sub>haloalkyl having 1 to 3 halogen atoms; C<sub>1</sub>-C<sub>2</sub>alkoxy-C<sub>1</sub>-C<sub>2</sub>alkyl; propenyl which is unsubstituted or substituted by 1 to 3 halogen atoms; propargyl; C<sub>3</sub>-C<sub>6</sub>cycloalkyl; C<sub>3</sub>-C<sub>6</sub>cycloalkylmethyl which is unsubstituted or substituted by 1 to 2 halogen atoms; cyano-C<sub>1</sub>-C<sub>2</sub>-alkyl;

C<sub>1</sub>-C<sub>2</sub>alkoxycarbonyl-C<sub>1</sub>-C<sub>2</sub>alkyl; phenyl-C<sub>1</sub>-C<sub>2</sub>alkyl which is unsubstituted or substituted by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano, nitro or C<sub>1</sub>-C<sub>2</sub>alkylenedioxy, it being possible for the phenyl group to be monosubstituted to disubstituted by identical or different substituents; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or pyridyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro.

24. A compound of the formula I according to claim 23, in which

A is an O atom,

R<sub>1</sub> is methyl,

R<sub>2</sub> is methyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, trifluoromethyl or trifluoromethoxy; and

R<sub>3</sub> is methyl.

25. A compound of the formula I according to claim 16, in which:



- 79 -

A is  $\text{NCH}_3$ ,

$\text{R}_1$  is methyl,

$\text{R}_2$  is methyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, trifluoromethyl or trifluoromethoxy;

$\text{R}_3$  is methyl; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or pyridyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro.

26. A compound of the formula I according to claim 1, in which:

X is an N atom;

Y is  $\text{OR}_{11}$ ;

$\text{R}_{11}$  is  $\text{C}_1$ - $\text{C}_4$ alkyl; and in which

A,  $\text{R}_1$ ,  $\text{R}_2$  and  $\text{R}_3$  are as defined in claim 1.

27. A compound of the formula I according to claim 26, in which:

A is an O atom;

$\text{R}_1$  and  $\text{R}_2$  are methyl;

$\text{R}_3$  is hydrogen;  $\text{C}_1$ - $\text{C}_4$ alkyl;  $\text{C}_1$ - $\text{C}_4$ haloalkyl having 1 to 3 halogen atoms;

$\text{C}_1$ - $\text{C}_2$ alkoxy- $\text{C}_1$ - $\text{C}_2$ alkyl; propenyl which is unsubstituted or substituted by 1 to 3 halogen atoms; propargyl;  $\text{C}_3$ - $\text{C}_6$ cycloalkyl;  $\text{C}_3$ - $\text{C}_6$ cycloalkylmethyl which is unsubstituted or substituted by 1 to 2 halogen atoms; cyano- $\text{C}_1$ - $\text{C}_2$ -alkyl;

$\text{C}_1$ - $\text{C}_2$ alkoxycarbonyl- $\text{C}_1$ - $\text{C}_2$ alkyl; phenyl- $\text{C}_1$ - $\text{C}_2$ alkyl which is unsubstituted or substituted by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano, nitro or  $\text{C}_1$ - $\text{C}_2$ alkylenedioxy; it being possible for the phenyl group to be monosubstituted to disubstituted by identical or different substituents; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or pyridyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro.

28. A compound of the formula I according to claim 27, in which

$\text{R}_3$  is hydrogen;  $\text{C}_1$ - $\text{C}_4$ alkyl or  $\text{C}_1$ - $\text{C}_4$ haloalkyl having 1 to 3 halogen atoms.

29. A compound of the formula I according to claim 1, in which:

- 80 -

X is an N atom;

Y is  $N(R_{12})R_{13}$ ;

$R_{12}$  and  $R_{13}$ , independently, are hydrogen atom or  $C_1$ - $C_4$ alkyl; and in which A,  $R_1$ ,  $R_2$  and  $R_3$  are as defined in claim 1.

30. A compound of the formula I according to claim 29, in which:

Y is  $NH_2$ ,  $N(CH_3)_2$  or  $NHC_2H_5$ ;

A is an O atom;

$R_1$  and  $R_2$  are methyl;

$R_3$  is hydrogen;  $C_1$ - $C_4$ alkyl;  $C_1$ - $C_4$ haloalkyl having 1 to 3 halogen atoms;

$C_1$ - $C_2$ alkoxy- $C_1$ - $C_2$ alkyl; propenyl which is unsubstituted or substituted by 1 to 3 halogen atoms; propargyl;  $C_3$ - $C_6$ cycloalkyl;  $C_3$ - $C_6$ cycloalkylmethyl which is unsubstituted or substituted by 1 to 2 halogen atoms; cyano- $C_1$ - $C_2$ alkyl;  $C_1$ - $C_2$ alkoxycarbonyl- $C_1$ - $C_2$ alkyl; phenyl- $C_1$ - $C_2$ alkyl which is unsubstituted or substituted by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano, nitro or  $C_1$ - $C_2$ alkylenedioxy, it being possible for the phenyl group to be monosubstituted to disubstituted by identical or different substituents; phenyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro; or pyridyl which is unsubstituted or monosubstituted to disubstituted, independently, by halogen, methyl, methoxy, halomethyl having 1 to 3 halogen atoms, cyano or nitro.

31. A compound of the formula I according to claim 30, in which

$R_3$  is hydrogen;  $C_1$ - $C_4$ alkyl or  $C_1$ - $C_4$ haloalkyl having 1 to 3 halogen atoms.

32. A compound of the formula I according to claims 1-31, in which the  $X=C$  double bond is in the E form.


33. A process for the preparation of a compound of the formula I according to claim 1, which comprises,

A) to prepare a compound of the formula I in which Y is  $N(R_{12})R_{13}$ : reacting a compound of the formula I in which Y is  $OR_{11}$  with  $HN(R_{12})R_{13}$  in a solvent at  $0^\circ C$  to  $40^\circ C$ ;

B) to prepare a compound of the formula I in which X, Y, A and  $R_1$ - $R_3$  are as defined for formula I (and where  $R_3$  is not hydrogen):


- 81 -

reacting an oxime of the general formula II



II

in which A and R<sub>1</sub>-R<sub>3</sub> are as defined above, with a benzyl derivative of the general formula




III

in which X and Y are as defined above and U is a leaving group, in an inert organic diluent, in the presence of a base and in the presence or absence of a phase transfer catalyst at 0°C to 50°;

C) to prepare a compound of the formula I in which A is oxygen and X, Y and R<sub>1</sub>-R<sub>3</sub> are as defined for formula I:

reacting an oxime of the general formula IV



IV

in which X, Y, R<sub>1</sub> and R<sub>2</sub> are as defined above, with a compound of the general formula V

U-R<sub>3</sub>


V,

- 82 -

in which  $R_3$  is as defined under formula I and U as defined under formula III (and where  $R_3$  is neither hydrogen nor phenyl nor pyridyl);

D) to prepare a compound of the formula IV in which X, Y,  $R_1$  and  $R_2$  are as defined for formula I:

reacting a ketone of the general formula VI




VI

in which X, Y,  $R_1$  and  $R_2$  are as defined above, with hydroxylamine or a salt thereof in a solvent at 0°C to 50°C;

E) to prepare a compound of the formula I in which A, X, Y and  $R_1$ - $R_3$  are as defined for formula I (and where  $R_3$  is not hydrogen):

reacting an enol or oxime of the general formula VII




VII

in which A, X, Y and  $R_1$ - $R_3$  are as defined above, with a methylating agent, for example methyl iodide, dimethyl sulfate or diazomethane in the presence of a base in a suitable solvent.

34. A process for the preparation of a compound of the formula VII


- 83 -



VII

in which A, X, Y and  $R_1$ - $R_3$  are as defined for formula I, which comprises,

A) to prepare a compound of the formula VII in which X is CH: reacting a phenylacetic acid derivative of the formula VIII




VIII

in which A, Y and  $R_1$ - $R_3$  are as defined above with a formate in the presence of a base;

B) to prepare a compound of the formula VII in which X is N: reacting a phenylacetic acid derivative of the formula VIII in which A, Y and  $R_1$ - $R_3$  are as defined above, with nitrous acid or a nitrite in the presence of a base;

C) to prepare a compound of the formula VII in which X is CH: reacting a keto ester of the formula IX



IX

in which A, Y and  $R_1$  -  $R_3$  are defined as above with methoxymethylenetriphenylphosphorane;

- 84 -

D) to prepare a compound of the formula VII in which X is N: reacting a keto ester of the formula IX in which A, Y and R<sub>1</sub>-R<sub>3</sub> are as defined above, with O-methylhydroxylamine or a salt thereof.

35. A composition for controlling pests, comprising, as active ingredient, a compound according to claim 1 together with a suitable carrier.

36. A composition according to claim 35 comprising, as active ingredient, a compound according to any one of claims 2-15.

37. A composition according to claim 35 comprising, as active ingredient, a compound according to any one of claims 16-25.

38. A composition according to claim 35 comprising, as active ingredient, a compound according to any one of claims 26-32.

39. A composition according to any one of claims 35-38, the pests being phytopathogenic microorganisms.

40. A composition according to claim 39, the microorganisms being fungi.

41. A composition according to any one of claims 35-38, the pests being insects or Acarina.

42. The use of a compound of the formula I according to claim 1 for controlling pests.

43. A method of controlling and preventing pests, which comprises applying a compound according to claim 1 to the pests or their environment.

44. A process according to claim 43, wherein a compound according to any one of claims 2-15 is applied.

45. A process according to claim 43, wherein a compound according to any one of claims 16-25 is applied.

- 85 -

46. A process according to claim 43, wherein a compound according to any one of claims 26-32 is applied.

47. A process according to any one of claims 43-46, the pests being phytopathogenic microorganisms.


48. A process according to claim 47, the microorganisms being fungi.

49. A process according to any one of claims 43-46, the pests being insects or Acarina.

50. A process according to any one of claims 43-46, wherein seed is treated.

51. Seed which has been treated according to claim 50.


52. A compound of the formula VII.



VII

in which A, X, Y and  $R_1$ - $R_3$  are as defined for formula I in claim 1.

53. A compound of the formula VIII




VIII

in which A, Y and  $R_1$ - $R_3$  are as defined for formula I in claim 1.

54. A compound of the formula IX

- 86 -



IX

in which A, Y and R<sub>1</sub>-R<sub>3</sub> are as defined for formula I in claim 1.



# INTERNATIONAL SEARCH REPORT

Int. Patent Application No  
PCT/EP 94/04318

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C07C251/60 A01N37/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07C A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,90 07493 (HOFFMANN-LA ROCHE AG) 12 July 1990  see example 60 see claims 1-4,9,10,14,15 ----	1,2,4,6, 7,9,10, 13,16, 21, 35-37, 39-45, 47-51
A	EP,A,0 472 300 (ICI) 26 February 1992 cited in the application see the whole document ----	1
A	EP,A,0 460 575 (CIBA-GEIGY AG) 11 December 1991 cited in the application see the whole document -----	1

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

22 February 1995

Date of mailing of the international search report

17. 03. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Goetz, G

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 94/04318

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO-A-9007493	12-07-90	AU-A-	4639089	01-08-90
		CA-A-	2005345	29-06-90
		EP-A-	0403618	27-12-90
		JP-T-	3503056	11-07-91
		NO-C-	173543	29-12-93
-----				
EP-A-0472300	26-02-92	US-A-	5346902	13-09-94
		US-A-	5238956	24-08-93
-----				
EP-A-0460575	11-12-91	AU-B-	633735	04-02-93
		AU-A-	7816791	12-12-91
		CA-A-	2043733	06-12-91
		JP-A-	4235953	25-08-92
-----				

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**